arXiv:1204.1741v2 [math.DS] 7 Dec 2012

DYNAMICS OF CONVEX COCOMPACT SUBGROUPS OF MAPPING CLASS GROUPS

ILYA GEKHTMAN

Abstract. For a convex cocompact subgroup $G < Mod(S)$, and points $x, y \in Teich(S)$ we obtain asymptotic formulas as $R \to \infty$ of $|B_R(x) \cap Gy|$ as well as the number of conjugacy classes of pseudo-Anosov elements in G of dilatation at most R. We do this by developing an analogue of Patterson-Sullivan theory for the action of G on PMF.

Contents

1. Statement of Results 1
1.1. Acknowledgments 4
2. Background on Teichmüller Theory 4
3. Fixing the Quasiconvexity Constants 8
4. Busemann Functions for the Teichmuller Metric 8
5. Conformal Densities for G 9
6. Patterson-Sullivan Construction of a Conformal density 10
7. Sectors and Closures 11
8. The Bowen Margulis Measure 14
9. Nonarithmeticity of the Length Spectrum 16
10. Ergodicity and Mixing of the Bowen Margulis Measure 18
11. Controlling the Multiple Zero Locus 23
12. Exact Asymptotics for Orbit Growth 28
13. Counting Closed Geodesics 40
References 43

1. Statement of Results

The study of the dynamics of the action of the mapping class group on Teichmüller space has long been influenced by analogy with the actions of discrete isometry groups of manifolds of negative curvature. Two properties of interest for a negatively curved manifold M are the

---

Date: 6 December 2012.

growth of orbits of $\pi_1(M)$ and the asymptotics as $R \to \infty$ of the number $n_M(R)$ of closed geodesics of length at most R. For M compact and negatively curved and $x, y \in M$, Margulis [12] showed in his 1970 thesis that

$$\lim_{R\to\infty} e^{-hR}|B_R(x) \cap \Gamma y| = \Lambda(x)\Lambda(y)$$

and

$$\lim_{R\to\infty} Re^{-hR} n_M(R) = 1$$

where Λ is a continuous function on M and h is the topological entropy of the geodesic flow. Roblin [22] generalized these results to $M = X/\Gamma$ any quotient of a contractible $CAT(-1)$ metric space by a geometrically finite group Γ. We prove an analogue of Roblin's result for certain subgroups of mapping class groups acting on Teichmüller space. Let S be a surface of genus $g \geq 2$. Let $Teich(S)$ be the associated Teichmüller space of isotopy classes of marked complex structures on S. Let $Mod(S) = Diff(S)/Diff_0(S)$ be the associated mapping class group, and let d_T denote the Teichmüller metric on $Teich(S)$. A subgroup $G < Mod(S)$ is called convex cocompact if its orbit in $Teich(S)$ is quasiconvex. Convex-cocompact subgroups of $Mod(S)$ were introduced by Farb and Mosher [2] and further developed by Kent and Leininger [14]. We prove:

Theorem 1.1. *Let S be a closed surface of genus $g \geq 2$. Let $G < Mod(S)$ be a convex cocompact subgroup containing a pseudo-Anosov element whose axis lies in the principal stratum. Let $x, y \in Teich(S)$ and $B_R(x)$ the ball of radius R about x in the Teichmüller metric. Let h be the exponent of convergence of G with respect to the Teichmüller metric. Then*

$$\lim_{R\to\infty} e^{-hR}|B_R(x) \cap \Gamma y| = \Lambda(x)\Lambda(y)$$

Theorem 1.2. *Let G be as in Theorem 1.1 Let $n_M(R)$ be the number of conjugacy classes of primitive pseudo-Anosov mapping classes in G of Teichmüller translation length at most R (this translation length is the logarithm of the dilatation of the pseudo-Anosov representative). Then*

$$\lim_{R\to\infty} Re^{-hR} n_M(R) = 1.$$

Like Margulis and Roblin, we prove the counting estimate by constructing a certain measure on the unit tangent bundle and prove it is mixing. We develop an analogue of Patterson-Sullivan theory for the action of $G < Mod(S)$ on Thurston's sphere PMF. We construct a unique G-conformal density $\nu_x, x \in Teich(S)$ supported on the limit set $\Lambda(G) \subset PMF$ of G, and scale the product measure $\nu_x \times \nu_x$ by a factor depending on the Busemann function to form a finite (in fact compactly supported) G invariant measure μ on the unit (co)tangent bundle $Q^1(S)$, which can be considered the analogue Bowen-Margulis measure in negative curvature. We prove

Theorem 1.3. *The measure μ associated with a convex cocompact subgroup $G < Mod(S)$ is mixing.*

Here G does not need to have a pseudo-Anosov whose axis lies in the principal stratum.

A difficulty faced in our setting is that $Teich(S)$ is not hyperbolic in any reasonable sense. It is neither $CAT(0)$ nor Gromov hyperbolic. Thurston proved that $Teich(S)$ has a natural $Mod(S)$ equivariant compactification by the sphere of projective measured foliations, but not every geodesic ray converges to a limit in PMF, rays with the same limit point are not necessarily asymptotic, and rays with different limit points may stay a bounded distance apart. Thurston's compactification coincides neither with the Gromov compactification (which is not Hausdorff) nor with the horofunction compactification (which contains PMF as a proper subset of smaller dimension). Nevertheless certain parts of $Teich(S)$ exhibit many aspects of hyperbolicity. Teichmüller geodesic triangles are δ thin as long as they lie over compact subsets of moduli space. Moreover, geodesic segments which spend a fixed proportion in quadratic differentials with no curves of short flat length behave like geodesics in $CAT(-1)$ metric spaces: namely, if two geodesics both pass within two balls of bounded radius lying far apart, they become very close somewhere in the middle. We use some arguments from ergodic theory to show that asymptotically, "most" geodesic segments are well-behaved in this sense and use techniques analogous to Roblin's to control the asymptotics of these well-behaved geodesics.

In order to prove mixing of μ we prove a certain nondegeneracy condition for the length spectrum of G, which is in our setting the measure of maximal entropy for the Teichmüller geodesic flow over $Teich(S)/G$.

Theorem 1.4. *Let $G < Mod(S)$ be a nonelementary subgroup. Then the logarithms of the dilatations of pseudo-Anosov elements of G generate a dense subgroup of $\mathbb{R}$.*

For subsemigroups of $SL_n\mathbb{R}$ acting irreducibly on $\mathbb{R}^n$ and containing a proximal element, an analogous result is proved by Guivarch and Urban in [23]. In variable negative curvature this question remains open. We prove Theorem 4 by using the piecewise linear structure of

MF given by train track coordinates to embedd a sub semigroup of G into $SL_n\mathbb{R}$ with the image satisfying the conditions of [23].

When G is the full mapping class group, analogues of Theorems 1.1 and 1.2 respectively were proved by Athreya-Bufetov-Eskin-Mirzakhani in [1] and Eskin-Mirzakhani in [5].

1.1. **Acknowledgments.** I would like to thank my advisor, Alex Eskin, for suggesting this problem to me and his constant advice, encouragement and insight. I would like to thank Howard Masur for teaching me a lot about Teichmüller theory and critical advice on many sections of this paper. I would like to thank Barbara Schapira for clearing up my misconceptions about Patterson-Sullivan theory and useful comments on an earlier draft of this paper. I would like to thank Alex Wright for useful comments on an earlier draft of this paper. I would like to thank Tam Nguyen Phan for teaching me a lot about various aspects of negative and nonpositive curvature. I would like to thank Frederic Paulin for an encouraging conversation about extending Patterson-Sullivan theory to the Teichmüller setting when I was beginning working on this project. I would like to thank Jayadev Athreya, Francoise Dal'bo, Benson Farb, Vaibhav Gadre, Roland Gunesch, and Joseph Maher for useful conversations.

2. Background on Teichmüller Theory

Let S be a closed surface of genus at least 2. Let $Mod(S)$ be the mapping class group of S. Let $Teich(S)$ be the space of marked complex structures on S up to isotopy. The space $Q(S)$ of quadratic differentials can be thought as a cotangent bundle of $Teich(S)$. A stratum of $Q(S)$ consists of all quadratic differentials whose zeros have the same combinatorial singularity type. The principal stratum consists of all quadratic differentials with simple zeros. Let $Q^1(S)$ be the space of area one quadratic differentials, which can be identified with the unit cotangent bundle to $Teich(S)$. Let $\pi : Q^1(S) \to Teich(S)$ be the projection. Let MF be the space of measured foliations on S and PMF its projectivization. For a quadratic differential q let $q^+, q^- \in MF$ denote its vertical and horizontal measured foliations respectively and $[q^+], [q^-] \in PMF$ its projective classes. Let $UE \subseteq PMF$ denote the projective classes of uniquely ergodic foliations. There is a compactification due to Thurston of $Teich(S)$ by PMF [7] obtained by embedding both into $\mathbb{R}^A$ where A is the set of isotopy classes of simple closed curves on S. Unless otherwise stated, the topology on

$Teich(S) \cup PMF$ in this paper comes from the Thurston compactification of $Teich(S)$. Given a basepoint in Teich(S), we can also compactify $Teich(S)$ by equivalence classes of geodesic rays through the point, called the Teichmüller compactification. Hubbard and Masur showed in [11] that there is a homeomorphism $Q(S) \to Teich(S) \times MF$ obtained by associating to a quadratic differential its projection in $Teich(S)$ and vertical (or horizontal) measured foliation. In [16] Masur shows:

Theorem 2.1. *If $q \in Q(S)$ with q^+ uniquely ergodic then $\pi(g_t q)$ converges to the projective class of q^+ in PMF.*

Theorem 2.2. *If $q_1, q_2 \in Q^1(S)$ and $q_1^+ = q_2^+$ is uniquely ergodic then $d_T(\pi(g_t q_1), \pi(g_t q_2)) \to 0$.*

The following is part of Masur's Two Boundaries Theorem [17]

Theorem 2.3. *The identity map on $Teich(S)$ extends to a homeomorphism between $T(S) \cup UE$ in the Teichmuller compactification and $T(S) \cup UE$ in the Thurston compactification.*

Theorem 2.4. *Let $x = x_0 \in Teich(S)$. Let $x_n \in Teich(S)$ be a sequence converging in the Thurston compactification to a uniquely ergodic $\eta \in PMF$. Then there exists a sequence of quadratic differentials $q_i \in S(x)$ and $t_i > 0$ such that $x_i = \pi(g_{t_i} q_i)$ and the q_i converge to $q \in S(x)$ such that $\eta = \lim_{t\to\infty} \pi(g_t q)$. Therefore for any fixed $m > 0$, the points $\pi(g_m q_i)$ converges to $\pi(g_m q_i)$.*

The following result of Klarreich is Prop 5.1 in [13].

Proposition 2.5. *Let F_1 and F_2 be topologically inequivalent minimal foliations. Let x_n and y_n be sequences in $Teich(S)$ converging to F_1 and F_2 respectively. Then the geodesic segments $[x_n, y_n]$ accumulate in the Teichmuller compactification to a set $s \subseteq Teich(S) \cup PMF$ such that $s \cap Teich(S)$ is a nonempty union of geodesics whose vertical and horizontal foliations are topologically equivalent to F_2 and F_1 respectively and $s \cap PMF$ consists of foliations that are equivalent to F_1 or F_2. If $x \in Teich(S)$ is fixed, then $[x, x_n]$ accumulate to a set $s \subseteq Teich(S) \cup PMF$ such that $s \cap Teich(S)$ is a union of geodesic rays based at x whose endpoints are topologically equivalent to F_1 and $s \cap PMF$ consists of foliations topologically equivalent to F_1. If F_1 and F_2 are topologically equivalent then $[x_n, y_n]$ converges to a subset of PMF each element of which is topologically equivalent to F_1.*

For uniquely ergodic foliations F_1 and F_2, the above proposition and Masur's two boundaries theorem implies the following.

Corollary 2.6. *$[x, x_n]$ converges uniformly on compact sets to $[x, [F_1])$ and $[x_n, y_n]$ converges to $([F_1], [F_2])$.*

The following corollary of Proposition 2.5 is proved in [14].

Corollary 2.7. *Let η_n, ζ_n in PMF converge to uniquely ergodic $\eta, \zeta \in PMF$. Then the accumulation points of $[\eta_n, \zeta_n]$ in the Thurston topology are contained in $\{\eta, \zeta\}$.*

Call a subgroup G of $Mod(S)$ non-elementary if it contains a pair of noncommuting pseudo-Anosovs. For a nonelementary subgroup G of $Mod(S)$, let $\Lambda(G)$ denote its limit set in PMF, the unique closed G invariant subset of PMF on which G acts minimally. The limit set is the closure of the set of stable foliations of pseudo-Anosov mapping classes in G. It is perfect and has empty interior provided it is not equal to PMF [15]. If any pair of points in $\Lambda(G)$ fill S, we define $WH(G)$ to be the union of all Teichmuller geodesics whose vertical and horizontal measured foliations have projective classes in $\Lambda(G)$. A subgroup G of $Mod(S)$ is called convex cocompact if some G-orbit in $Teich(S)$ is quasiconvex. The following properties of convex cocompact subgroups of $Mod(S)$ are proved in [2] and [14].

Theorem 2.8.
- *Every G orbit is quasi-convex.*
- *The weak hull $WH(G)$ is defined and G acts cocompactly on $WH(G)$.*
- *Every limit point η of G is conical, that is for $x \in Teich(S)$ there is some $D > 0$ such that the ray $[x, \eta)$ has infinite intersection with D neighborhood of Gx .*
- *G acts cocompactly on $WH(G) \cup \Lambda(G)$.*
- *$WH(G)$ is contained in the ϵ-thick part of $Teich(S)$ and A-quasiconvex for some $\epsilon > 0$ and $A > 0$.*
- *$WH(G) \cap \Lambda(G)$ is closed in $Teich(S) \cap PMF$.*
- *Every point of $\Lambda(G)$ is uniquely ergodic.*
- *G contains a finite index subgroup all of whose nonidentity elements are pseudo-Anosov.*

From now on, let G denote a nonelementary convex cocompact subgroup of $Mod(S)$.

Lemma 2.9. *If G is nonelementary convex cocompact, $\Lambda(G)$ coincides with $\overline{Gx} \cap PMF$ for any $x \in Teich(S)$.*

Proof. Let $\eta \in \Lambda(G)$ be an the attracting point of a pseudo-Anosov $g \in G$. Then $g^n x \to \eta$ so $\eta \in \overline{Gx} \cap PMF$. Since the fixed points of pseudo-Anosovs of G are dense in $\Lambda(G)$, we have $\Lambda(G) \subseteq \overline{Gx} \cap PMF$ (this holds for any nonelementary $G \subseteq Mod(S)$. For the other direction, it is proved by McCarthy and Papadopoulos in [18] that $\overline{Gx} \cap PMF$ is contained in

$$Z\Lambda(G) = \{\lambda \in PMF | \exists \beta \in \Lambda(G) : i(\lambda, \beta) = 0\}.$$

Since every point of $\Lambda(G)$ is uniquely ergodic, $Z\Lambda(G) = \Lambda(G)$. $\square$

Let $Teich_\epsilon(S)$ denote the ϵ thick part of $Teich(S)$, the set of hyperbolic structures on S where no closed curve has hyperbolic length less than ϵ. The following property of Teichmüller geodesics, indicative of hyperbolicity in the thick part, is proved by Rafi [21].

Lemma 2.10. *For each $A > 0$ and $\epsilon > 0$ there exists a constant $D > 0$ such that for points $x, x', y, y' \in Teich_\epsilon(S)$ with $d_T(x, x') \leq A$ and $d_T(y, y') \leq A$ the geodesic segments $[x, y]$ and $[x', y']$ D-fellow travel in a parametrized fashion, and for $\eta \in PMF$ such that $[x, \eta)$ and $[x', \eta)$ are contained in $Teich_\epsilon(S)$, the geodesic rays $[x, \eta)$ and $[x', \eta)$ D-fellow travel in a parametrized fashion.*

For a subset W of a metric space and $A > 0$ let $N_A W$ denote the A neighborhood of W.

Corollary 2.11. *Let $G \leq Mod(S)$ be convex cocompact. For every $C > 0$ there exists an $C' > 0$ such that every geodesic with endpoints in $\Lambda(G) \cup N_C WH(G)$ is contained entirely in $N_{C'} WH(G)$.*

Proof. Let $A > 0$ be such that $WH(G) \cup \Lambda(G)$ is A-quasiconvex. Let $\epsilon > 0$ be such that $N_C WH(G) \subseteq Teich_\epsilon(S)$. (Such an ϵ exists since G acts cocompactly on $WH(G)$ and therefore on $N_C WH(G)$). Then for each $x, y \in N_C WH(G)$ there are $x', y' \in WH(G)$ with $d_T(x, x') \leq C$ and $d_T(y, y') \leq C$. Since x, x', y, y' lie in $Teich_\epsilon(S)$ and $d_T(x, x') \leq C$ and $d_T(y, y') \leq C$, it follows from Rafi's theorem that the geodesic segments $[x, y]$ and $[x', y']$ D-fellow travel. Since $WH(G)$ is A-quasiconvex, $[x', y']$ is contained in $N_{A+C} WH(G)$ and so $[x, y]$ is contained in $N_{A+C+D} WH(G)$. The proof when one of x, y lies in $\Lambda(G)$ is similar. $\square$

The following is proved in [19].

Proposition 2.12. *For every $\epsilon > 0$ there exists an $\delta > 0$ such that any triangle with vertices in $Teich(S) \cup PMF$ and sides contained in $Teich_\epsilon(S)$ is δ thin- each side is contained in a δ neighborhood of the other two.*

From the proof of [14], Theorem 4.4 we also have the following.

Proposition 2.13. *For each $\epsilon > 0$ there exists a $K > 0$ with the following property. Suppose $x, y, z \in Teich(S) \cup PMF$ form a triangle with sides contained in the ϵ thick part of $Teich(S)$. Let $P \in [x, y]$ minimize the distance between $[x, y]$ and z. Then $[x, P] \cup [P, z]$ lies in a K neighborhood of $[x, z]$.*

3. Fixing the Quasiconvexity Constants

Existence of the following constants is guaranteed by the above remarks. Fix $A > 0$.
Let $A'' > 0$ be such that any geodesic between two points in

$$N_A WH(\Lambda(G)) \cup \Lambda(G)$$

is contained in

$$N_{A''} \cup WH(\Lambda(G)) \cup \Lambda(G)$$

Let $A' > 0$ be such that any geodesic between two points in

$$N_{A''} WH(\Lambda(G)) \cup \Lambda(G)$$

is contained in

$$N_{A'} \cup WH(\Lambda(G)) \cup \Lambda(G)$$

Let $\epsilon > 0$ be such that $N_{A'} WH(\Lambda(G)) \subseteq Teich_\epsilon(S)$. Let $K > 0$ be large enough so that any triangle in Teichmüller space with sides contained in $Teich_\epsilon(S)$ is K thin, satisfies Proposition 2.13, and also large enough such that the shadow $pr_\eta B_K(x)$ contains an open set intersecting $\Lambda(G)$ for every $x \in N_{A'} WH(G)$ and $\eta \in \Lambda(G)$ (see section 7 below).

4. Busemann Functions for the Teichmuller Metric

If $x, y \in Teich(S)$, $\alpha \in MF$ uniquely ergodic and and $z_n \to [\alpha]$ in the Thurston compactification, then Miyachi [20] showed

$$d_T(x, z_n) - d_T(y, z_n) \to \frac{1}{2} \log \frac{Ext_\alpha(x)}{Ext_\alpha(y)}.$$

In particular, the limit $\beta_{[\alpha]}(x, y) = d_T(x, z_n) - d_T(y, z_n)$ exists and varies continuously with $[\alpha] \in UE$. This gives a continuous extension of the cocycle

$$\beta_z(x, y) = d_T(x, z) - d_T(y, z)$$

to $Teich(S) \cup UE$.

For $\zeta, \eta \in MF$ uniquely ergodicand $x, w, z \in Teich(S)$ we can also define

$$\rho_x(z, w) = d(x, z) + d(x, w) - d(z, w)$$

$$\rho_x(z,[\zeta]) = \rho_x([\zeta],z) = \lim_{w\to[\zeta]} d(x,z)+d(x,w)-d(z,w) = d(x,z)+\beta_{[\zeta]}(x,z)$$

$$\rho_x([\zeta],[\eta]) = \lim_{(z,w)\to([\zeta],[\eta])} d(x,z)+d(x,w)-d(w,z) = \beta_{[\zeta]}(x,u)+\beta_{[\eta]}(x,u)$$

$$= \frac{1}{2}\log\frac{Ext_x\zeta Ext_x\eta}{i(\eta,\zeta)^2}$$

where $u \in Teich(S)$ is any point on the Teichmuller geodesic defined by ζ,η. The function ρ is continuous in $x \in Teich(S)$ and $\zeta,\eta \in UE$. It can be considered as an analogue in our setting of the Gromov product.

5. Conformal Densities for G

A conformal density for G is a family $\{\nu_x | x \in Teich(S)\}$ of borel measures on PMF, each supported on $\Lambda(G)$ satisfying
(1)

$$\gamma * \nu_x = \nu_{\gamma x}$$

for all $x \in Teich(S)$ and $\gamma \in G$ and
(2) For all $x, y \in Teich(S)$ ν_x and ν_y are absolutely continuous and satisfy

$$\frac{d\nu_x}{d\nu_y}(\alpha) = exp(\delta(G)\beta_\alpha(x,y).$$

Note, by the G-invariance of the Busemann cocycle if condition (2) is satisfied, it suffices to check condition (1) at a single x.

Proposition 5.1. *A conformal density ν_x for a nonelementary convex cocompact $G < Mod(S)$ has full support on $\Lambda(G)$ and has no atoms.*

Proof. Suppose $U \subseteq PMF$ is open and $U \cap \Lambda(G) \neq \emptyset$ but $\nu_x(U) = 0$. Since the limit set is the closure of the set of stable (or unstable) laminations of pseudo-Anosov mapping classes in G, there is some pseudo Anosov $\gamma \in G$ with axis l with repelling fixed point $l^- \in U$. Then for each $n > 0$,

$$\nu_x(\gamma^n U) = \nu_{\gamma^{-n}x}(U) = 0$$

since ν_x and $\nu_{\gamma^{-n}x}$ are absolutely continuous. Note

$$\bigcup_{n>0} \gamma^n U = PMF \setminus l^+$$

By countable subadditivity of the measure, ν_x is concentrated on the single point l^+. However, since G is not elementary, there is some $h \in G$ with $hl^+ \neq l^+$, and by absolute continuity we must also have

$$\nu_x(hl^+) = \nu_{h^{-1}x}(l^+) \neq 0$$

giving a contradiction. Thus we have proved that ν_x has full support on $\Lambda(G)$. Now, suppose ν_x has an atom $\eta \in \Lambda(G)$, say of mass r. By

$[KL1]$ every limit point η of G is conical, that is there exists a $D > 0$ such that the D neighborhood of the geodesic $[x, \eta)$ intersects the orbit Gx infinitely many times. Let $\gamma_n \in G$ be such a sequence. Then by the triangle inequality, $\beta_\eta(\gamma_n x, x) \to \infty$. Then,

$$\nu_x(\gamma_n^{-1}\eta) = \nu_{\gamma_n x}(\eta) = exp(\delta(G)\beta_\eta(\gamma_n x, x))\nu_x(\eta) \to \infty$$

contradicting the finiteness of ν_x. □

6. Patterson-Sullivan Construction of a Conformal density

Let δ_G be the exponent of convergence of G. For $s > \delta_G$ and $x, y \in Teich(S)$ let

$$f_s(x, y) = \sum_{\gamma \in G} \exp(-sd(x, \gamma(y))).$$

Fix $x \in Teich(S)$. Now let

$$\nu_{x,s} = f_s(x, x)^{-1} \sum_{\gamma \in G} \exp(-sd(x, \gamma(x)))\delta_{\gamma x}$$

where δ_p denotes the dirac measure at p. Now, consider a weak-* limit ν_x of the $\nu_{x,s}$ as $s \to \delta_G$. It is a probability measure on

$$\overline{Teich(S)} = Teich(S) \cup PMF.$$

Assume first that the Poincare series diverges at δ_G. Then $f(s) \to \infty$, so by discreteness of G, ν_x gives zero measure to compact subsets of $Teich(S)$, and thus must be supported on PMF.

Furthermore, since each $\nu_{x,s}$ is supported on the G orbit of x, it follows that ν_x is supported on its closure, so it must be supported on $\Lambda(G)$. For any other $y \in Teich(S)$ define

$$d\nu_y(\alpha) = \exp(\delta(G)\beta_\alpha(x, y))d\nu_x.$$

Since

$\beta_\alpha(x, z) = \beta_\alpha(x, y) + \beta_\alpha(y, z)$

we have that

$$\frac{d\nu_y}{d\nu_z}(\alpha) = \exp(\delta(G)\beta_\alpha(z, y)).$$

Now, we show that this gives a conformal density. Indeed, for any $g \in G$ we have,

$$g\nu_{gx,s}(z) = exp(-s\beta_z(gx, x))\nu_{x,s}(z)$$

where β is the Busemann cocycle, and taking limits as $s \to \delta$ we get

$$gd\nu_x(\alpha) = \exp(-\delta(G)\beta_\alpha(gx, x))d\nu_x$$

so we indeed have a conformal density.

Now, suppose the Poincare series converges at $\delta(G)$ (this case turns out to be vacuous, but the construction of a conformal density is required to show it). There exists a slowly growing function h on $\mathbb{R}$ such that

$$\sum_{\gamma \in G} h(d(x, \gamma x)) exp(-sd(x, \gamma(y)))$$

diverges at $s = \delta(G)$ but converges for $s < \delta(G)$. We then set

$$f_s(x, x) = \sum_{\gamma \in G} h(d(x, \gamma x) exp(-sd(x, \gamma(x)))$$

and carry out the construction as before. The existence of an appropriate function h is guaranteed by application of the following result of [26] to the Radon measure

$$\sum_{\gamma \in \Gamma} D_{d(x, \gamma x)}.$$

Lemma 6.1. *Let λ be a Radon measure on $\mathbb{R}_+$, such that the Laplace transform of λ*

$$\int_{\mathbb{R}_+} e^{-st} d\lambda(t)$$

has critical exponent $\delta \in \mathbb{R}$. Then there exists a nondecreasing function $h : \mathbb{R}_+ \to \mathbb{R}_+$ such that

$$\int_{\mathbb{R}_+} h(t) e^{-\delta t} d\lambda(t) = \infty$$

and for every $\epsilon > 0$ there exists $t_0 \geq 0$ such that, for any $u \geq 0$ and $t \geq t_0$, one has

$$h(u + t) \leq e^{\epsilon t} h(t)$$

In particular the Laplace transform of $h\lambda$ has critical exponent δ.

7. Sectors and Closures

For $x, y \in T(S)$ and $\eta \in PMF(S)$ let $pr_\eta(x) \in PMF(S)$ denote the vertical projective measured foliation of the quadratic differential $q \in S(x)$ with horizontal projective measured foliation η and $pr_y(x) \in PMF(S)$ the vertical projective measured foliation of the geodesic segment from x to y.

Lemma 7.1. *The function $pr_{\cdot}(*)$ is continuous on $(Teich(S) \cup UE) \times Teich(S)$.*

Proof. The continuity on $Teich(S) \times Teich(S)$ and $PMF \times Teich(S)$ follows from the fact that the map $S(x) \to PMF$ is a homeomorphism for $x \in Teich(S)$. If $x_n \to \eta$ uniquely ergodic, then $[x_n, x]$ converges to $(\eta, x]$. Let $y_n \in [x_n, x]$ be at distance one from x. Then $y_n \to y = \gamma_{x,\eta}(1)$ so $pr_{x_n}(x) = pr_{y_n}(x) \to pr_y(x) = pr_\eta(x)$. □

For $x \in Teich(S)$ and $U \subseteq PMF$ let $Sect_x(U)$ be the set of all $y \in Teich(S)$ with $pr_x(y) \in U$ For any $r > 0$ let

$$C_r^+(x,U) = N_r \bigcup_{z \in B_r(x)} Sect_x(U)$$

and

$$C_r^-(x,U) = \left\{ y \in Teich(S) | B(y,r) \subseteq \bigcap_{z \in B(x,r)} Sect_z(U) \right\}$$

Lemma 7.2. *For any $r > 0$, $D > 0$ and $U \subset PMF$ open the closure of $C_r^\pm(x,U) \cap N_D WH(G)$ in the Thurston compactification is*

$$(C_r^\pm(x, \overline{U}^{PMF}) \cap N_D WH(G)) \cup (\overline{U}^{PMF} \cap \Lambda(G))$$

and the closure of $Sect_x(U) \cap N_D WH(G)$ in the Thurston compactification is

$$Sect_x(\overline{U}^{PMF}) \cap N_D WH(G)) \cup (\overline{U}^{PMF} \cap \Lambda(G))$$

Proof. Since *pr* is continuous, $Sect_x \overline{U}^{PMF}$ and $C_r^\pm(x, \overline{U}^{PMF})$ are the closures in $Teich(S)$ of $Sect_x(U)$ and $C_r^\pm(x,U)$ respectively. Since $N_D WH(G)$ is closed in $Teich(S)$ we have that $Sect_x \overline{U}^{PMF} \cap N_D WH(G)$ and $C_r^\pm(x, \overline{U}^{PMF}) \cap N_D WH(G)$ are the closures in $Teich(S)$ of $Sect_x(U) \cap N_D WH(G)$ and $C_r^\pm(x,U) \cap N_D WH(G)$ respectively.

Now let $x_n \in C_r^+(x,U) \cap N_D WH(G)$ converge to some $\eta \in PMF$. Since $x_n \in N_D WH(G)$ we have $\eta \in \Lambda(G)$. By definition of C_r^+ there exists a sequence $y_n \in B_r(x_n) \cap Sect_x U$. Since η is uniquely ergodic we have $y_n \to \eta$. Now let $z_n \in [x, x_n]$ with $d(z_n, x) = 1$ and $z \in [x, \eta)$ with $d(z_n, x) = 1$. Since η is uniquely ergodic we have $z_n \to z$ and by continuity of *pr* we have $pr_x z_n \to pr_x z = \eta$. By definition $pr_x z_n \in U$ so $\eta \in \overline{U}$.

Now, suppose $\eta \in \Lambda(G) \cap U$. Then $\pi g_t q_{x,\eta} \to \eta$ as $t \to \infty$. Let $D' > 0$ be such that $[x, \alpha) \subset N_{D'} WH(G)$ for all $\alpha \in \Lambda(G)$. Then $w_t = \pi g_t q_{x,\eta} \in N_{D'} WH(G)$ for all t. Let $v_t \in WH(G) \cap B_{D'}(w_t)$. As $\eta \in UE$ we have $z_t \to \eta$. We claim $B_{r+D}(w_t \subset Sect_x(U)$ for large enough t, whence it will follow that $v_t \in C_r^-(x,U)$ for large enough t. Indeed, otherwise, letting $V \subset PMF$ be an open neighborhood of η

with closure in U there is a sequence $t_n \to \infty$ and $p_n \in B_{r+D'}(w_{t_n}) \setminus Sect_xU$. Since $\eta \in UE$ we have $p_n \to \eta$ and thus $pr_xp_n \to \eta$. Thus $pr_xp_n \in U$ for large enough n since U is open in PMF, contradicting our assumption. So $v_t \in WH(G) \cap C_r^-(x,U)$ converges to η.

□

Corollary 7.3. *If $U, V \subset PMF$ with $\overline{U} \subset V^o$ then $(C_r^+(x,U) \cap N_DWH(G)) \setminus C_r^-(y,V)$ has compact closure in $Teich(S)$.*

Corollary 7.4. *If V is open in $Teich(S) \cup PMF$ and $U \subset PMF$ with $\overline{U} \subset V \cap PMF$ then $C_r^+(x,U) \setminus V$ has compact closure in $Teich(S)$. If X is closed in $Teich(S) \cup PMF$ and $U \subset PMF$ is open in PMF with $X \cap PMF \subset U$ then $X \setminus C_r^-(x,U)$ has compact closure in $Teich(S)$.*

Let $V \subset PMF$ be open with $\overline{U} \subset V$.

Lemma 7.5. *For any geodesic l with endpoints in $\Lambda(G) \setminus V$, $l \cap Sect_x(U)$ is contained in a compact subset of $Teich(S)$. (In particular l spends only a finite amount of time in $Sect_xU$).*

Proof. Suppose $l \cap Sect_x(U)$ is not contained in a compact subset of $Teich(S)$. Then (for a correct choice of orientation of the geodesic) there exist $s_n \to \infty$ such that $p_n = l_{s_n} \in Sect_xU$. Note, $l_{s_n} \to l^+ \in \Lambda(G) \setminus V$, which is uniquely ergodic. Let $q_n \in S(x)$ and $t_n > 0$ be such that $p_n = \pi(g_{t_n}q_n)$. Note, as $p_n \in Sect_x(U)$ we have $[q_n+] \in \overline{U}$. Then $q_n \to q \in S(x)$ with

$$l^+ = [q^+] = \lim_{t \to \infty} \pi(g_tq)$$

But since the map

$$S(x) \to PMF$$
$$q \to [q^+]$$

is a homeomorphism, $[q^+] \in \overline{U}$ contradicting that $l^+ \in \Lambda(G) \setminus V$. □

Lemma 7.6. *There exists a $T > 0$ such that any geodesic l with $l^+, l^- \in \Lambda(G) \setminus V$ and $d(x,l) \geq T$ is disjoint from $Sect_xU$.*

Proof. Suppose not. Then there is a sequence l_n of geodesics with $l_n^+, l_n^- \in \Lambda(G) \setminus V$ and $d(x,l) > n$ and $p_n \in l_n \cap Sect_xU$. Passing to a subsequence, we have either that (l_n^+, l_n^-) converges to either a pair of distinct points (η, ζ) in $\Lambda(G)$ or a single point $\eta \in \Lambda(G)$. In the first case, we would have l_n converge in the Hausdorff topology on $Teich(S) \cup PMF$ to the geodesic l between η and ζ, so $d(l_n,x) \to d(l,x)$ which would contradict $d(x,l_n) \to \infty$. Thus, we have l_n^+ and l_n^- converging to the same $\eta \in \Lambda(G) \setminus V$. Then, we have l_n converging to

η and thus $p_n \to \eta$. Let $q_n \in S(x)$ be such that $p_n = \pi(g_{t_n} q_n)$. Note, as $p_n \in Sect_x(U)$ we have $[q_n^+] \in \overline{U}$. Then $q_n \to q \in S(x)$ with

$$\eta = [q^+] = \lim_{t\to\infty} \pi(g_t q)$$

But $[q^+] \in \overline{U}$ contradicting that $\eta \in \Lambda(G) \setminus V$. ☐

Lemma 7.7. *There exists a $D > 0$ such that for every $\eta, \zeta \in \Lambda(G) \setminus V$ the geodesic between η, ζ spends at most time D in $Sect(U)$.*

Proof. Suppose the contrary. Then there exist a sequence of geodesics l_n with both endpoints l_n^+, l_n^- outside of V such that l_n spends time at least n in $Sect(U)$. We may pass to a subsequence such that one of the following holds: either (l_n^+, l_n^-) converges to a pair of distinct endpoints (η, ζ) in $\Lambda(G) \setminus V$ or both converge to the same $\eta \in \Lambda(G) \setminus V$. Suppose the first case. Then, the geodesics l_n converge uniformly on compact sets to the geodesic l with endpoints η, ζ. Thus, this geodesic must spend an infinite amount of time inside $Sect(U)$, which is impossible if both of its endpoints are outside of U. Now suppose both endpoints of l_n converge to the same $\eta \in \Lambda(G) \setminus V$. Then, l_n converges to η. Let $p_n \in l_n \cap Sect_x(U)$. Let $q_n \in S(x)$ be such that $p_n = \pi(g_{t_n} q_n)$. Note, as $p_n \in Sect_x(U)$ we have $[q_n+] \in \overline{U}$. Then $q_n \to q \in S(x)$ with

$$\eta = [q^+] = \lim_{t\to\infty} \pi(g_t q)$$

But $[q_n+] \in \overline{U}$ contradicting that $\eta \in \Lambda(G) \setminus V$. ☐

8. The Bowen Margulis Measure

Let $x \in Teich(S)$. Define a measure $\widetilde{\mu}$ on $\Lambda(G) \times \Lambda(G)$ by

$$d\tilde{\mu}(\eta, \zeta) = \exp(\delta(G)\rho_x(\eta, \zeta)) d\nu_x(\eta) d\nu_x(\zeta)$$

Note, every distinct pair of points in $\Lambda(G)$ give a Teichmuller geodesic in $WH(G)$ so we can consider $\tilde{\mu}$ as a measure on $Q^1(S)$, invariant under the Teichmüller geodesic flow g_t and supported on $WH(G)$. By continuity of ρ on pairs of points in the limit set, $\tilde{\mu}$ is locally finite.

Lemma 8.1. *The measure $\tilde{\mu}$ is G invariant.*

Proof. This follows from the G equivariance property of conformal densities. ☐

Thus $\tilde{\mu}$ descends to a measure μ on $Q^1(S)/G$. Since G acts cocompactly on $WH(G)$, it follows that μ is compactly supported, and therefore finite. We call μ the Bowen-Margulis measure. By projecting μ to $Teich(S)$ we obtain a measure on $Teich(S)$, supported on

$WH(G)$. For $q_0 \in Q^1(S)$ let the strong stable (unstable) leaf associated to q_0, denoted by $W^{ss}(q_0)$ (resp $W^{su}(q_0)$) be the set of elements of $Q^1(S)$ with the same vertical (resp horizontal) measured foliation as q_0. Let the weak stable (unstable) leaf associated to q_0, denoted by $W^s(q_0)$ (resp $W^u(q_0)$) be the set of elements of $Q^1(S)$ with the same vertical (resp horizontal) projective measured foliation as q_0.

The map sending each quadratic differential to its horizontal projective measured foliation restricts to a map

$$P_-^{q_0} : W^{ss}(q_0) \to PMF$$

that is a homeomorphism between $W^{ss}(q_0)$ and $PMF \setminus V(q_0^+)$ where for a measured foliation α, $V(\alpha)$ consists of the foliations θ such that $i(\alpha,\beta)+i(\theta,\beta)=0$ for some $\beta \in PMF$. In particular, if q_0^+ is uniquely ergodic, $P_-^{q_0}$ is a homeomorphism between $W^{ss}(q_0)$ and $PMF \setminus q_0^+$. We can define a locally finite measure on $W^{ss}(q_0)$, denoted by $\mu^{\tilde{ss}}_{q_0}$ by pulling back the Patterson-Sullivan measure on PMF and scaling by the Busemann function:

$$d\mu^{\tilde{ss}}_{q_0}(v) = \exp(-\delta\beta_{[v^-]}(x, \pi(v)))d\nu_x([v^-])$$

Here, $x \in Teich(S)$ is any basepoint and μ^{ss} is independent of x. Similarly, we can define measure on strong unstable horospheres $W^{su}(q_0)$ by

$$d\mu^{\tilde{su}}_{q_0}(v) = \exp(-\delta\beta_{[v+]}(x, \pi(v)))d\nu_x([v^+])$$

We can also define measures on weak horospheres by integrating the measures on strong horospheres with respect to geodesic arclength.

$$\widetilde{\mu^{s}_{q_0}} = \widetilde{\mu^{ss}_{q_0}} dg_t$$

$$\widetilde{\mu^{u}_{q_0}} = \widetilde{\mu^{su}_{q_0}} dg_t$$

These project modulo Γ to measures $\mu^{su}_{q_0}$ and $\mu^u_{q_0}$, $\mu^s_{q_0}$ and $\mu^s_{q_0}$.

Note, whenever q_0^+ is uniquely ergodic, there is a map

$$h^{ss}_{q_0} : Q^1(S) \setminus W^u(-q_0) \to W^{ss}(q_0)$$

with

$$h^s_{q_0}(q) = W^u(q) \cap W^{ss}(q_0)$$

which has one to one restrictions to any unstable horosphere. When q^+ is uniquely ergodic, $h^{ss}_{q_0}$ restricts to a homeomorphism between $S(x) \setminus p$ and $W^{ss}(q_0)$ where $p^+ = q_0^+$. We can also define a map

$$h^s_{q_0} : Q^1(S) \setminus W^u(-q_0) \to W^s(q_0)$$

with

$$h^s_{q_0}(q) = W^{su}(q) \cap W^s(q_0)$$

Similarly define maps h^u and h^{su}

Lemma 8.2. *For any q_1, q_2 with q_i^+ uniquely ergodic, the restriction of $h^s_{q_1}$ to $W^s(q_2)$ takes $\widetilde{\mu^s_{q_2}}$ to $\widetilde{\mu^s_{q_1}}$*

We can now define, for a fixed $q_0 \in Q^1(S)$, a measure $d\tilde{\mu}_{q_0}$ on $Q^1(S)$ by

$$d\tilde{\mu}_{q_0} = dg_w(g_t w) d\mu^{ss}_v(w) d\mu^{uu}_{q_0}(v)$$

ie by integrating over $g_t w$ with geodesic arclength dg_w, then integrating over all w in $W^{ss}(v)$ with respect toμ^{ss}_v, finally integrating over all v in $W^{su}(q_0)$ with respect to $\mu^{su}_{q_0}$.

Lemma 8.3. *μ_{q_0} is independent of q_0 and coincides with the Bowen-Margulis measure $\tilde{\mu}$.*

Proof. This follows from the fact that ν_x is supported on the uniquely ergodic part of PMF and that $\beta_{[v^+]}(x, \pi(w)) = \beta_{[v^+]}(x, \pi(v))$ whenever $q \in W^{ss}(v)$ and v^+ uniquely ergodic. ☐

9. Nonarithmeticity of the Length Spectrum

In this section we prove

Theorem 9.1. *Let $G < Mod(S)$ be a nonelementary subgroup. The the logarithms of the dilatations of pseudo-Anosov elements of G generate a dense subgroup of $\mathbb{R}$.*

For a train track τ, let $W_\tau \cong \mathbb{R}^{6g-6}$ be the vector space of weights on the branches of τ satisfying the switch condition. Let $V_\tau \subset W_\tau$ be the open cone assigning positive measure to each branch. Each element of V_τ corresponds to a measured foliation. Let $\phi_\tau : V_\tau \to MF$ be this correspondence. Let U_τ be the image of ϕ_τ in MF and let $\psi_\tau : U_\tau \to V_\tau$ be the inverse of ϕ_τ.

Lemma 9.2. *Let $\gamma \in Mod(S)$ be a pseudo-Anosov such that γ^+ is carried by the interior of the maximal recurrent train track τ and γ^- is not carried by τ. Then for large enough n we have $\gamma^n \tau$ is carried by τ and γ^n acts linearly on W_τ by a positive matrix whose largest eigenvalue is the dilatation $\lambda(\gamma)$ of γ.*

Let $G < Mod(S)$ be nonelementary.

Let τ be a maximal recurrent train track and $\gamma_1, \gamma_2 \in G$ independent pseudo-Anosovs such that γ_i^+ are carried by τ, and γ_i^- are not carried by τ; replacing γ_i by high enough iterates we can assume that they preserve U_τ. Let $\Gamma \subset G$ be the semigroup freely generated by the γ_i. Then each $\gamma \in \Gamma$ preserves U_τ and acts linearly on $W\tau$ by a positive matrix whose largest eigenvalue is the dilatation $\lambda(\gamma)$ of γ. Moreover, since the mapping class group preserves Thurston's symplectic form on $Mod(S)$ we have that each $\gamma \in \Gamma$ acts on W_τ by a symplectic matrix.

Note, if A represents the action of $\gamma \in G$ then E_A cannot have any elements of V_τ, ie cannot have any vectors with all entries nonnegative. Indeed, if $v \in E_A$ then $[A^n v]$ does not converge to $[v_A] \in PW_\tau$. However, if $v \in V_\tau$ then $v = \psi_\tau \alpha$ for some measured foliation α with $[\alpha] \neq [\gamma^-]$. Thus, $[\gamma^n \alpha] \to [\gamma^+]$ and hence

$$[A^n v] = [A^n \psi_\tau \alpha] = [\psi_\tau(\gamma^n \alpha)] \to [\psi_\tau \gamma^+] = v_A$$

In particular we obtain that if independent elements $\gamma_1, \gamma_2 \in \Gamma$ act on W_τ by matrices A and B respectively then $\Gamma v_B \cap E_A = \emptyset$.

Thus, it suffices to prove the following result about linear semigroup actions on projective space. For a proximal element $A \in SL_n\mathbb{R}$ let v_A be a dominant eigenvector with corresponding eigenvalue $\lambda(A)$ and E_A the direct sum of complementary eigenspaces. We need

Theorem 9.3. *Let Γ be a semigroup of $SL_n\mathbb{R}$ every element of which is proximal. Suppose for any $A, B \in \Gamma$ we have $\Gamma v_B \cap E_A = \emptyset$. Then the logarithms of maximal eigenvalues of matrices in Γ generate a dense subgroup of $\mathbb{R}$.*

Let the limit set of Γ, denoted by L_Γ be the closure in $P\mathbb{R}^n$ of

$$\{[v_A] : A \in \Gamma\}$$

Let $\widetilde{L_\Gamma}$ be the preimage of L_Γ in $\mathbb{R}^n$.

Lemma 9.4. *L_Γ is Γ invariant.*

Proof. Suppose $A, B \in \Gamma$. We need to show that $[Av_B] \in L_\Gamma$. Consider

$$u = \lim_{n\to\infty} \frac{1}{||B||^n} B^n \in M_N(\mathbb{R})$$

Then u is a projection onto $\mathbb{R}v_B$ with $ker u = E_B$ and $u(v_B) = v_B$ By assumption, $Av_B \notin E_B = ker u = ker Au$. Thus, Au is a multiple of a projection onto $\mathbb{R}Av_B$. Note,

$$Au = \lim_{n\to\infty} \frac{1}{||B||^n} AB^n$$

so $[v_{AB^n}] \to [Av_B]$. Hence, $[Av_B] \in L_\Gamma$. □

Lemma 9.5. *Any Γ invariant subspace W is either contained in $\bigcap_{A\in\Gamma} E_A$ or contains v_A for all $A \in \Gamma$.*

Proof. Suppose $\Gamma W = W$, and $B \in \Gamma$ with $v_B \notin \Gamma$. Then for any $v \notin E_B$ we have $\lim_{k\to\infty}[B^k v] = [v_B] \notin [W]$. However, for any $v \in W$ we have $B^k v \in W$ and as $[W]$ is closed, any limit point of $\{[B^k v] : k \in \mathbb{N}\}$

is in $[W]$. Thus, if $v_B \notin W$ then $W \subset E_B$. In particular, since for any $A \in \Gamma$ we have $v_A \notin E_B$ we have that W does not contain v_A for any $A \in \Gamma$ and is thus contained in E_A for all $A \in \Gamma$. $\square$

Let $W_\Gamma = \oplus_{A\in\Gamma}\mathbb{R}v_A$ be the smallest subspace of $\mathbb{R}^n$ containing $\widetilde{L_\Gamma}$. Since Γ preserves L_Γ, it preserves W_Γ. Let U_Γ be a maximal proper Γ invariant subspace of W_Γ.

Lemma 9.6. *Γ acts irreducibly on $V_\Gamma = W_\Gamma/U_\Gamma$, and each $A \in \Gamma$ has the same largest eigenvalue in this action as in the action on $\mathbb{R}^n$.*

Proof. The irreducibility follows from maximality of U_Γ. Note, by Lemma 10.4 we have $U_\Gamma \subset \bigcap_{A\in\Gamma} E_A$ does not contain any v_A for $A \in \Gamma$ so $v_A + U_\Gamma$ is a dominant eigenvector for A with eigenvalue $\lambda(A)$. $\square$

Lemma 9.7. *For any independent $A, B \in \Gamma$ there is an integer $M > 0$ such that $\Gamma_M = sg(A^M, B^M)$ acts strongly irreducibly on $V_{\Gamma_M} = W_{\Gamma_M}/U_{\Gamma_M}$.*

Proof. Now, consider independent $A, B \in \Gamma$. Since there is no infinite nested sequence of finite dimensional subspaces, there exists an $N > 0$ and subspaces $U \subset W$ such that $W_{\Gamma_M} = W$, $U_{\Gamma_M} = U$ and $V_{\Gamma_M} = W/U = V$ for all $M \geq N$. We know that $\Gamma_N = sg(A^N, B^N)$ acts irreducibly on V and consequently so does the free group $G_n =< A^N, B^N >$ generated by A^N, B^N. Suppose the Γ_N action on V is not strongly irreducible. Let $V_1, ..., V_n \subset V$ be a minimal collection of subspaces of V such that their union is preserved by Γ_N. Note, A^N, B^N permute $V_1, ..., V_N$ so there exists an $K > 0$ such that $A^{MK}V_i = V_i$ and $B^{MK}V_i = V_i$ for each i. In particular, $\Gamma_{MK} = sg(A^{MK}, B^{MK})$ preserves the proper nontrivial subspace $V_1 \subset V = W$, contradicting the fact that Γ_{MK} must act on V irreducibly. $\square$

The following is proved in [23], Proposition 4.9.

Lemma 9.8. *Let $\Gamma < GL_n\mathbb{R}$ be a semigroup acting irreducibly on $\mathbb{R}^n$ and containing a proximal element. Then the logarithms of maximal eigenvalues of proximal elements of Γ generate a dense subgroup of $\mathbb{R}$.*

Theorem 9.1 follows from Lemmas 9.6 and 9.7.

10. Ergodicity and Mixing of the Bowen Margulis Measure

Theorem 10.1. *The geodesic flow g_t on $Q^1(S)/G$ is ergodic with respect to the Bowen-Margulis measure μ.*

Proof. For $f \in C_c(Q^1(S)/G)$ continuous with compact support, consider the forward and backward Birkhoff averages:

$$f^+(q) = \lim\sup_{T\to\infty} \frac{1}{T}\int_0^T g_t q dt$$

and

$$f^-(q) = \lim\sup_{T\to\infty} \frac{1}{T}\int_0^T g_{-t} q dt$$

By the Birkhoff ergodic theorem these are finite and equal for almost every $q \in Q^1(S)/G$. Moreover, it is clear that f^+ and f^- are invariant under geodesic flow. Furthermore, f^+ is invariant along $W^{ss}(q)$ whenever q^+ is uniquely ergodic, and f^- is invariant along $W^{su}(q)$ whenever q^- is uniquely ergodic. Suppose the measure is not ergodic. Then there exists some $f \in C_c(Q^1(S)/G)$ such that f^+ is NOT almost everywhere constant. Let C_1, C_2 be disjoint sets whose union is $\mathbb{R}$ such that $D_i = (f^+)^{-1}C_i$ has positive measure. Note, by Fubini's theorem and the product structure of the measure μ there exists q_0 with $q_0- \in \Lambda(G)$ and a set $A \subseteq W^{su}(q_0)$ of full $\mu^{su}_{q_0}$ measure such that $f^+(v) = f^-(v)$ and $v^+ \in \Lambda(v)$ for all $v \in A$. Furthermore there are sets $A_i(q_0) \subseteq A$ of positive $\mu^{su}_{q_0}$ measure such that for all $v \in A_i(q_0)$ $W^s(v)$ intersects D_i in a set of positive $\mu^{ss}_v \times dt$ measure. Note, since the D_i are g_t invariant and f^+ is constant along $W^{ss}(v)$, it follows that $W^s(v) \subseteq D_i$ for $v \in A_i(q_0)$. In particular, $A_i(q_0) \subseteq D_i$. However, as $f^+ = f^-$ on A this implies that $f^-(A_i)(q_0) \subseteq C_i$. However, as q_0- is uniquely ergodic, f^- is constant on $W^{su}(q_0)$ contradicting that the C_i are disjoint. ☐

It follows that G acts ergodically on PMF with $\nu_x \times \nu_x$.

Theorem 10.2. *The geodesic flow g_t on $Q^1(S)/G$ is mixing with respect to the Bowen-Margulis measure μ.*

Our argument is modelled on Babbillot's argument in [24] where an analogous result was proved for general quasi-product measures on manifolds of pinched negative curvature. The following result from unitary representation theory is proved in [24].

Theorem 10.3. *Let $(X, B, m, (T_t)_{t\in A})$ be a measure preserving dynamical system where (X, B) is a Borel space, m a Borel measure on X and T an action an action of a locally compact second countable abelian group on X by m preserving transformations. Let $f \in L^2(X, m)$, and if m is finite assume also $\int_X f dm = 0$. Then, if $f \circ T_a$ does not converge weakly to 0 as $a \to \infty$ in A, there exist a sequence $s_n$going to infinity* *in* A *and a non-constant function* $\psi \in L^2(X, m)$ *such that* $f \circ T_{s_n}$ *and* $f \circ T_{-s_n}$ *both converge to* ψ*.*

Suppose μ is not mixing. Then there is a continuous G invariant function f with $supp(f)/G$ compact such that

$$\int_{Q^1} f d\tilde{\mu} = 0$$

and $f \circ g_t$ does not converge weakly to zero. Let $s_n \to \infty$ and nonconstant ψ be such that $\int_{Q^1} f dm = 0$ and both $f \circ g_{s_n}$ and $f \circ g_{-s_n}$ converge weakly to ψ. By the Banach-Saks theorem, there exists a subsequence t_n of s_n such that the Cesaro averages

$$A_N = \frac{1}{N} \sum_{n=1}^{N} f \circ g_{t_n}$$

and

$$A_{-N} = \frac{1}{N} \sum_{n=1}^{N} f \circ g_{-t_n}$$

converge almost surely to ψ. We first smooth out ψ by considering the function $v \to \int_0^c \psi(g_s v) ds$. Choosing small enough c guarantees that this function remains non-constant, and it is moreover the limit of the corresponding Cesaro averages of the smoothing of f. By abuse of notation, we continue to call the new functions f and ψ. Now, there exists a set E_0 of full μ measure in $\Lambda(G) \times \Lambda(G)$ such that for each v on a geodesic with endpoints in E_0, the function $t \to \psi(g_t(v))$ is well defined and continuous. Consider the closed (a priori possibly trivial) subgroup $\mathbb{R}(q)$ of $\mathbb{R}$ given by the periods of $t \mapsto \psi(g_t q)$. It is clearly flow invariant, and thus gives a measurable map from E_0 into the set of closed subgroups of $\mathbb{R}$. By ergodicity of $\nu_x \times \nu_x$ it must be constant almost everywhere on E_0. Suppose this subgroup is $\mathbb{R}$. Then ψ would be g_t invariant, and thus pass to a flow invariant function on Q^1/G, which is not almost-everywhere constant. However, this contradicts the ergodicity of μ. Thus the subgroup in question must be cyclic. Say it equals $k\mathbb{Z}$ on a full measure set $E_1 \subseteq E_0$. Let

$$\psi^+ = \limsup A_N$$

and

$$\psi^- = \limsup A_{-N}$$

By Fubini's theorem, there is a set $E_2 \subseteq E_1$ be of full measure and such that $\psi^+ = \psi^- = \psi$ everywhere along every geodesic in E_2.

Now, let E^- be the set of $\lambda \in \Lambda(G)$ such that for ν_x almost everywhere α, $(\lambda, \alpha) \in E_2$ and similarly let E^+ be the set of $\lambda \in \Lambda(G)$

such that for ν almost everywhere α, $(\alpha, \lambda) \in E_2$. Again, by Fubini's theorem, $E = E_2 \cap (E^+ \times E^-)$ has full measure.

Now, let $\eta_1, \eta_2, \zeta_1, \zeta_2 \in \Lambda(G)$. Choose $p_0 \in (\eta_1, \zeta_1)$, $p_1 \in (\zeta_1, \eta_2)$, $p_2 \in (\eta_2, \zeta_2)$, $p_3 \in (\zeta_2, \eta_1)$ and $p_4 \in (\eta_1, \zeta_1)$ such that p_i and p_{i+1} are on the same horosphere. We claim that the distance $\tau(\eta_1, \eta_2, \zeta_1, \zeta_2)$ between p_0 and p_4 depends only on the η_i and ζ_i and is thus independent of the position of p_0 on its geodesic. It will follow that this distance is a period of $t \mapsto f(g_t q_0)$ where $q_0 \in S(p_0)$ with $[q_0^+] = \zeta_1$ and thus is contained in $k\mathbb{Z}$ for ν_x almost every $\eta_1, \eta_2, \zeta_1, \zeta_2 \in \Lambda(G)$. Indeed, let H_{η_i} and H_{ζ_i} be horospheres centered at η_i and ζ_i respectively for $i = 1, 2$. Let D_{ij} be the signed distance between the intersections of (η_i, ζ_j) with H_{η_i} and H_{ζ_j}, with the sign convention chosen in such a way that D_{ij} is positive if H_{η_i} and H_{ζ_j} are disjoint. Then since the geodesic flow takes horospheres to horospheres, the quantity $D_{1,1} + D_{2,2} - D_{1,2} - D_{2,1}$ is independent of the specific choice of horospheres. Moreover if the horospheres are chosen in such a way that H_{ζ_1} contains p_0, H_{η_2} passes through $p_1 = H_{\zeta_1} \cap (\zeta_1, \eta_2)$, H_{ζ_2} passes through $p_2 = H_{\eta_2} \cap (\eta_2, \zeta_2)$, and $H(\eta_1)$ passes through $p_3 = H_{\zeta_2} \cap (\eta_1, \zeta_2)$, then $D_{1,1} + D_{2,2} - D_{1,2} - D_{2,1}$ reduces to the signed distance $D_{1,1}$ between p_0 and $p_4 = H_{\eta_1} \cap (\eta_1, \zeta_1)$. Thus, $\tau(\eta_1, \eta_2, \zeta_1, \zeta_2) = D_{1,1} + D_{2,2} - D_{1,2} - D_{2,1}$ is well-defined and continuous on quadruples of points in $\Lambda(G)$. We call it the cross ratio of the four points in PMF, or the cross ratio of the geodesics (η_1, ζ_1) and (η_2, ζ_2).

Proposition 10.4.

$$\tau(\eta_1, \eta_2, \zeta_1, \zeta_2) = \lim_{n\to\infty} d(x_1^n, y_1^n) + d(x_2^n, y_2^n) - d(x_1^n, y_2^n) - d(y_1^n, x_2^n)$$

where $x_i^n, y_i^n \in Teich(S)$ *with* $x_i^n \to \eta_i$, $y_i^n \to \zeta_i$.

Proof. Let H_{η_i} and H_{ζ_i} be pairwise disjoint horospheres through η_i and ζ_i respectively, and let M_0 be the complement in $Teich(S)$ of the corresponding horoballs. The intersections of the geodesics

$$(\eta_1, \zeta_1), (\zeta_1, \eta_2), (\eta_2, y_2), (y_2, \eta_1)$$

with M_0 consist of disjoint segments I_j of length $d_j, j = 1, 2, 3, 4$ respectively. Note the number

$$\tau' = d_1 + d_2 - d_3 - d_4$$

does not depend on the specific choice of horosphere and by continuity of the Busemann function on $\Lambda(G)$ depends continuously on the $\eta_i, \zeta_i \in \Lambda(G)$. We claim that it is equal to $\tau(\eta_1, \eta_2, \zeta_1, \zeta_2)$. Indeed, suppose $x_1^n, x_2^n, y_1^n, y_2^n$ are points in $Teich(S)$ converging to η_i and ζ_i respectively. The segments $[x_1^n, y_1^n] \to (\eta_1, \zeta_1)$ $[x_1^n, y_2^n] \to (\eta_1, \zeta_2)$

$[x_2^n, y_1^n] \to (\eta_2, \zeta_1)$ $[x_2^n, y_2^n] \to (\eta_2, \zeta_1)$ uniformly on compact sets. In particular, their intersection with M_0 contains four segments I_j^n which converge toward the I_j. Thus, to prove that $\tau = \tau'$ it suffices to show that the contribution to

$$d(x_1^n, y_1^n) + d(x_2^n, y_2^n) - d(x_1^n, y_2^n) - d(y_1^n, x_2^n)$$

of the parts of $[x_i^n, y_j^n]$ which are contained in the complement of M_0 goes to zero. By symmetry it suffices to show that if p_1^n is the intersection of $[x_1^n, y_1^n]$ with H_{η_1}, p_2^n is the intersection of $[x_1^n, y_2^n]$ with H_{η_1}, then

$$d(p_1^n, x_1^n) - d(p_2^n, x_1^n) \to 0$$

Note, as $n \to \infty$ we have

$$p_1^n \to p_1 = H_{\eta_1} \cap (\eta_1, \zeta_1)$$

and

$$p_2^n \to p_2 = H_{\eta_1} \cap (\eta_1, \zeta_2)$$

and $\beta_{x_1^n}$ converges to β_{η_1} uniformly on compact sets. Thus

$$\beta_{x_1^n}(p_1^n, p_2^n) = d(p_1^n, x_1^n) - d(p_2^n, x_1^n) \to \beta_{\eta_1}(p_1, p_2)$$

which is zero since p_1, p_2 lie on the same horosphere based at η_1 □

From the expressions of the Busemann functions in terms of extremal length, we in fact find

$$\tau([\alpha_1], [\alpha_2], [\beta_1], [\beta_2]) = \frac{1}{2} \log \frac{i(\alpha_1, \beta_1) i(\alpha_2, \beta_2)}{i(\alpha_1, \beta_2) i(\alpha_2, \beta_1)}$$

for any α_i, β_i uniquely ergodic. Note, τ defines a continuous function on quadruples of points in $Teich(S) \cup \Lambda(G)$ From this formula we obtain

Corollary 10.5. *For any pseudo-Anosov $g \in Mod(S)$, with fixed points $\eta_1, \eta_2 \in \Lambda(G)$ the translation distance of λ is twice $\tau(\eta_1, \eta_2, \beta, g\beta)$ where β is any uniquely ergodic point in PMF distinct from the η_i.*

As noted above, $\tau(\eta_1, \eta_2, \zeta_1, \zeta_2)$ is a period of $t \mapsto f(g_t q_0)$ where $q_0 \in S(p_0)$ with $[q_0^+] = \zeta_1$ and so $\tau(\eta_1, \eta_2, \zeta_1, \zeta_2) \in k\mathbb{Z}$ for ν_x^4 almost every $\eta_1, \eta_2, \zeta_1, \zeta_2 \in \Lambda(G)$. By continuity of the cross ratio and the fact that ν has full support on $\Lambda(G)$, it follows that $\tau(\eta_1, \eta_2, \zeta_1, \zeta_2) \in k\mathbb{Z}$ for ν_x^4 for every $\eta_1, \eta_2, \zeta_1, \zeta_2 \in \Lambda(G)$. But this implies that the translation length of every element of G is in $k\mathbb{Z}$ contradicting Theorem 9.1.

11. Controlling the Multiple Zero Locus

By ergodicity, μ gives full mass to a single stratum. In the remainder of the paper, we will assume that this is the principal stratum. Here is a sufficient condition.

Proposition 11.1. *If G contains a pseudo-Anosov element with axis lying in the principal stratum, then μ gives full weight to the principal stratum.*

Proof. By proposition 5.1, ν_x has full support on $\Lambda(G)$ and thus μ has full support on $Q^1WH(G)$. Thus, any open set $U \subset Q^1(S)$ intersecting any geodesic with endpoints in $\Lambda(G) \times \Lambda(G)$ has positive μ measure. If $\gamma \in G$ has axis g_γ in the principal stratum then a point $p \in g_\gamma$ has a neighborhood $U \subset Q^1(S)$ that is also contained in the principal stratum. Thus, the principal stratum has positive μ measure and by ergodicity of μ on $Q^1(S)/G$ it has full measure. ☐

In this section, we show that the contribution to orbit growth of the multiple zero locus and thin parts of the principal stratum is asymptotically negligible. For a subset $P \subset Q^1(S)$, $c \in (0,1)$ and $x \in Teich(S)$ let $B_R(x,P,c)$ the set of points $y \in Teich(S)$ with $d_T(x,y) \leq R$ and the segment $[x,y]$ spending a proportion at most c of the time in P.

For $x, y \in Teich(S)$ let $N_G(x,y,P,R,c)$ denote the number of $\gamma \in G$ such that $d(x,\gamma y) \leq R$ and $[x,\gamma y]$ spending a proportion at most c of the time in P. Specifically, we prove:

Theorem 11.2. *For each $x, y \in Teich(S)$ and $\epsilon > 0$ there exists a closed subset $P' \subset Q^1(S)$ disjoint from the multiple zero locus such that*

$$\limsup_{R\to\infty} N_G(x,y,P',R,1/3)/e^{\delta R} \leq \epsilon$$

In order to prove this we will show:

Theorem 11.3. *For each $\epsilon > 0$ there exists a closed subset $P \subset Q^1(S)$ disjoint from the multiple zero locus such that*

$$\limsup_{R\to\infty} e^{-\delta R} m(B_R(x,P,1/2)) \leq \epsilon$$

We first conclude Theorem 11.2 from Theorem 11.3. We will a lemma of Eskin and Mirzakhani from [5]

Lemma 11.4 ([5], Lemma 5.4)**.** *Suppose $K \subset M_g$ is compact. Given $s > 0$, there exists constants L_0 depending on s and K, and c_0 depending only on K with the following property. If $\gamma : [0,L] \to Q^1(S)$ is a geodesic segment (parametrized by arclength) with endpoints above K*

and $L > L_0$*,* $\widehat{\gamma} : [0, L'] \to Q^1(S)$ *is the geodesic segment connecting* $p_1, p_2 \in Teich(S)$ *such that* $d_T(p_1, \pi(\gamma(0))) < c_0$*,* $d_T(p_2, \pi(\gamma(L))) < c_0$*, and*

$$|\{s \in [0, L] | l_{min}(\gamma(t)) \geq s\}| > \frac{L}{2}$$

then

$$|\{s \in [0, L'] | l_{min}(\widehat{\gamma}(t)) \geq s/4\}| > \frac{L'}{3}$$

From this, we obtain:

Lemma 11.5. *Let* $K \subset Teich(S)$ *be a compact subset, and* P *a closed subset of the principal stratum. Then there exists a closed subset* P' *of the principal stratum containing* P *in its interior and an* $R_0 > 0$ *such that for* $y_1, y_2 \in K$ *and* $x \in Mod(S)K$ *with* $d_T(x, y_i) > R_0$*, if* $[x, y_1]$ *spends a proportion at most* $1/3$ *in* P' *then* $[x, y_2]$ *spends a proportion at most* $1/2$ *in* P*.*

In particular we have:

Lemma 11.6. *For any* $K \subset Teich(S)$ *compact and* P *a closed subset of the principal stratum, there exists a closed subset* P' *of the principal stratum containing* P *in its interior and an* $R_0 > 0$ *such that for any* $x \in GK$*,* $y_1, y_2 \in K$ *and* $R \geq R_0$ *we have*

$$N_G(x, y_1, P', R, 1/3) \leq N_G(x, y_2, P, R + diam(K)).$$

Proof of Theorem 11.2 assuming Theorem 11.3. Let $x \in Teich(S)$ and $P \subset Q^1(S)$ be a subset of the principal stratum be such that the conclusion of Theorem 11.3 holds with $\epsilon/e^{\delta diam(K)}$ in place of ϵ, ie

$$m(B_R(x, P, 1/2)) \leq \frac{\epsilon}{e^{\delta diam(K)}} e^{\delta R}$$

for all large enough R.

Let $K \subset Teich(S)$ be compact and contain both x and fundamental domain for the action of G on $WH(G)$.

Lemma 11.7.

$$m(B_R(x, P, 1/2)) = \int_{y \in K} N_G(x, y, P, R, 1/2).$$

Proof. Note,

$$m(B_R(x, P, 1/2)) = \sum_{g \in G} \int_{y \in gK} \chi_{B_R(x,P,1/2)}(y) dm(y) = \int_{y \in K} \sum_{g \in G} \chi_{B_R(x,P,1/2)}(gy) dm(y) =$$

$$\int_{y \in K} N_G(x, y, P, R, 1/2).$$

□

Note, by Lemma 11.6 there exists a closed subset P' of the principal stratum containing P in its interior and a $R_0 > 0$ such that if $R > R_0$, then

$$N_G(x, y_1, P', R, 1/3) \leq N_G(x, y_2, P, R + diam(K))$$

for any $y_1, y_2 \in K$. Moreover, by Lemma 11.7 with with $R + diam(K)$ in place of R, for each large enough R there exists a $y_2 \in K$ such that

$$N_G(x, y_2, P, R{+}diam(K)), 1/2) \leq m(B_{R+diam(K)}(x, P, 1/2)) \leq \frac{\epsilon}{e^{\delta diam(K)}} e^{\delta(R+diam(K))} = \epsilon e^{\delta R}$$

completing the proof. □

We now consider the following measure on MF: Note the space of strong stable (or unstable) horospheres based at uniquely egodic points can be identified with uniquely ergodic points of MF. Indeed, let $o \in Teich(S)$ be a basepoint. If $\eta \in MF$ with $Ext_o \eta = 1$ then $t\eta$ is identified with the horosphere $H(t\eta) = H(t, [\eta])$ based at η such that $\beta_{[\eta]}(o, z) = t$ for each $z = z_{t,[\eta]} \in H$ (ie $Ext_z \eta = e^{2t}$). For $A \subset MF$ so that $[A] \subset PMF$ let

$$\lambda(A) = \int_{[\eta] \in [A]} \int_{t: H(t,[\eta]) \in A} e^{\delta t} d\nu_o([\eta]) = \int_{[\eta] \in [A], Ext_o \eta = 1} \int_{t\eta \in A} e^{\delta t} dt d\nu_o([\eta])$$

Lemma 11.8. *The measure λ does not depend on choice of basepoint $o \in Teich(S)$ and is G invariant. It has support precisely on foliations projecting to points on $\Lambda(G)$. Moreover for all $U \subset Q^1(S)$*

$$\lambda(\eta^+(g_t U)) = e^{\delta t} \lambda(\eta^+ U)$$

and

$$\mu(U) = \int_{\eta \in MF} \mu^{ss}(A \cap H(\eta)) d\lambda(\eta)$$

Proof. The independence of basepoint and G invariance follows because the ν_o form a conformal density, the other properties are immediate from the definition. □

Denote

$$\overline{\lambda}(U) = \lambda(Cone(U))$$

where $Cone(U)$ is the union of segments from the origin in MF to points of U. For $W \subset Q^1(S)$ and $s > 0$ let $W(s)$ denote the set of $q \in Q^1(S)$ such that there exists $q' \subset W$ on the same leaf of W^{su} as q such that $d_H(q, q') < s$. For a subset $A \subset Teich(S)$ let $A(r) = Nbhd_r(A)$ denote the r neighborhood of A in the Teichmüller metric.

Lemma 11.9. *Let $K \subset Teich(S)$ be a fundamental domain for the action of G on $WH(G)$. Let $h > 0$. Then there is a $C(h) > 0$ depending only on K and h such that for all $U \subset Q^1WH(G) \cap \pi^{-1}K$ and all $t > 0$ letting $W_t = g_tU$ we have*

$$m(Nbhd_2\pi(W_t)) \leq C(h)\overline{\lambda}(\eta^+W_t(h)).$$

Proof. Let $h_0 = h_0(K, h)$ be a small constant to be specified later. We can decompose U into pieces U_α such that each piece is within Hodge distance $h_0/2$ of a single unstable leaf. The minimal number of such pieces can be bounded by a constant depending only on K by the compactness of K and equivalence of the Euclidean and Hodge metrics over compact sets so, we may assume without loss of generality that U is within Hodge distance $h_0/2$ of a single unstable leaf. Also, as in [ABEM, Lemma 4.1], we can assume without loss of generality that U has $W^{su} \times W^s$ product structure. Pick a maximal $\Delta \subset \pi(W_t)$ with $d_T(x, y) = 1$ for any distinct $x, y \in \Delta$. Note by compactness of K and G equiariance of m there is a constant $C(K)$ depending only on K such that

$$m(B(X, 3)) \leq C(K)$$

for all $X \in GK$ Then

$$Nbhd_2\pi(W_t) \subset \bigcup_{X\in\Delta} B_T(X, 3),$$

and hence

$$m(Nbhd_2\pi(W_t)) \leq \sum_{X\in\Delta} (B_T(X, 3)) \leq |\Delta|C(K)$$

Now, let $\Delta' \subset W_t$ be a set containing one element of $\pi^{-1}(X) \cap Q^1WH(G)$ for each $X \in \Delta$. Let $B^{su}_E(q, r)$ denote the elements of $W^{su}(q)$ within euclidean distance r of q. As shown in [1], Lemma 4.1 for h_0 small enough we can pick h_2 depending only on K such that the $\eta^+(B^{su}(q, h_2)$ for distinct $q \in \Delta'$ are disjoint viewed as subsets of PMF. By equivalence of Hodge and Euclidean metrics there is a $h_3 \in (0, h_2)$ such that whenever $q \in \pi^{-1}K$ with $q' \in B^{su}_E(q, h_3)$ we have $d_H(q, q') \leq h$. For each $q \in \Delta'$ consider

$$H(q) = \eta^+(B^{su}_E(q, h_3)) \subset \eta^+(W(h)).$$

These are pairwise disjoint. Note, since ν_x has full support on $\Lambda(G)$, $\overline{\lambda}(H(q)) > 0$ for all $q \in Q^1WH(G)$. Thus, as $\overline{\lambda}$ is G equivariant and G acts cocompactly on $Q^1WH(G)$, there is a $c = c(K, h)$ such that

$$\overline{\lambda}(H(q)) \geq c$$

for all $q \in Q^1WH(G)$. Thus

$$\overline{\lambda}(\eta^+(W(h))) \geq \sum_{q \in \Delta'} \overline{\lambda}(\eta^+(H(q))) \geq c|\Delta|.$$

This completes the proof. $\square$

Now, let $P_1 \subset Q^1(S)/G$ be compact (in our application P_1 we will be a subset of the principal stratum of almost full μ measure) and define $P_3 \subset P_2 \subset P_1$ and $\delta \in (0, 1]$ such that if $q \in P_i$ and $d_H(q, q') \leq c_H h$ then $q' \subset P_{i-1}$ where c_H is the nonexpansion constant of the modified Hodge norm over P_1. By choosing h small enough we can assume $\mu(P_3) > 1/2$. For $T_0 > 0$ let $U'_i = U'_i(T_0)$ be the set of $q \in Q^1WH(G)$ such that there exists $T > T_0$ so that $g_t q$ is in the complement of P_i for at least half of $t \in [0, T]$. By definition $U'_1 \subset U'_2 \subset U'_3$ and by the Birkhoff ergodic theorem, for every $\theta > 0$ there is a $T_0 > 0$ such that $\mu(U'_3) < \theta$. Let $U_i = p^{-1}U_i \cap \pi^{-1}K$.

Lemma 11.10. *In the above notation, for all* $t > 0$

$$m(Nbhd_2(\pi(g_t U_1))) \leq C(h) e^{\delta t} \overline{\lambda}(\eta^+(U_2))$$

and for any $\epsilon > 0$ *it is possible to choose* T_0 *such that for all* $t > T_0$

$$m(Nbhd_2(\pi(g_t U_1))) \leq \epsilon e^{\delta t}.$$

Proof. Let $W = g_t U$. As shown in the proof of [1], Lemma 4.2 we have $W(h) \subset g_t U_2$. Now, we can apply Lemma 11.9 to W and use the fact that

$$\overline{\lambda}((\eta^+ g_t U)) = e^{\delta t} \overline{\lambda}(\eta^+ U)$$

to get the first claim. Moreover, as shown in the proof of [ABEM, Lemma 4.2], if $q \in U_2$ and $q' \in Q^1WH(G)$ is on the same strong stable leaf as q with $d_H(q, q') < h$ then $q' \in U_3$. By compactness of $Q^1WH(G)/G$ and G equivariance of μ^{ss}, there is a $c > 0$ such that $\mu^{ss}(B^{ss}(q)) > c$ for all $q \in Q^1WH(G) \cap p^{-1}K$. Therefore, by the product structure of μ, $\lambda(U_2) \leq C_1(h)\mu(U_3)$ where $C_1(h)$ depends only on h. Hence, choosing a large enough T_0 the second claim of the lemma follows. $\square$

Proof of Theorem 11.3. In the above notation, let P_1 be chosen disjoint from the multiple zero locus. Let T_0, U_1, U_1 be as in the proof of Lemma 11.10. Let $K \subset Teich(S)$ be a fundamental domain for the action of G on $WH(G)$ (so m is supported on $GK = WH(G)$). Then for $R > T_0$ and $x \in K$ we have

$$B_R(X, P_1) \cap GK \subset \bigcup_{0 \leq t \leq R} \pi(g_t U_1) \cap GK \subset \bigcup_{n=0}^{\lfloor R \rfloor} \bigcup_{n \leq t \leq n+1} \pi(g_t U_1) \cap GK$$

Then,

$$m(Nbhd_1 B_R(X,P)) \leq \sum_{n=0}^{\lfloor R \rfloor} m(Nbhd_2(\pi(g_n U_1)) \cap GK) \leq C\epsilon \sum_{n=0}^{\lfloor R \rfloor} e^{\delta n}$$

by Lemma 11.4. This completes the proof for $x \in WH(G)$. $\square$

12. Exact Asymptotics for Orbit Growth

The goal of this section is to prove the part of Theorem 1.1 concerning orbit growth. For $r > 0$, $x \in Teich(S)$ and $A \subseteq PMF$ recall

$$C_r^+(x,A) = N_r \bigcup_{z \in B_r(x)} Sect_z(A)$$

and

$$C_r^-(x,A) = \left\{ y \in Teich(S) | B(y,r) \subseteq \bigcap_{z \in B(x,r)} Sect_z(A) \right\}$$

For $t > 0$ and $x, y \in Teich(S)$ define a measure

$$d\nu_{x,y}^t = \delta ||\mu|| e^{-\delta t} \sum_{d(x,\gamma y) \leq t} D_{\gamma x} \otimes D_{\gamma^{-1} y}$$

Proposition 12.1. *Let $c > 0$, $x, y \in Teich(S)$ and $\eta_0, \zeta_0 \in PMF$ be such that there exist $\eta_0^*, \zeta_0^* \in \Lambda(G)$ with $x \in (\eta_0, \eta_0^*)$, $y \in (\zeta_0, \zeta_0^*)$. Then there exist open neighborhoods V and W in PMF of η_0 and ζ_0 respectively such that for all borel $A \subseteq V$ and $B \subseteq W$ with nonempty interior, as $T \to \infty$ we have*

$$\limsup \nu_{x,y}^T (C_1^-(x,A) \times C_1^-(y,B)) \leq e^c \nu_x(A) \nu_y(B)$$

and

$$\liminf \nu_{x,y}^T (C_1^+(x,A) \times C_1^+(y,B)) \geq e^{-c} \nu_x(A) \nu_y(B)$$

Proof. If η_0 is not in $\Lambda(G)$ then we can choose a neighborhood U of η_0 in PMF with $\nu_x(U) = 0$ and $W = PMF$ so that both sides of the desired equation are 0. Similarly if ζ_0 is not in $\Lambda(G)$. Assume therefore that $\eta_0, \zeta_0 \in \Lambda(G)$. The argument is modelled on Roblin's Theorem 4.1.1 in [22], where an analogous result is proved for manifolds of pinched negative curvature. For $\eta, \zeta \in PMF$ filling and $z \in Teich(S)$, let $z_{\eta,\zeta}$ be the quadratic differential with projective vertical and horizontal measured foliations ζ and η respectively and such that $\pi(z_{\eta,\zeta})$ lies at minimal distance from z. For $z \in Teich(S)$, $r > 0$ and $A \subseteq PMF$ let

$$K^+(z,r,A) = \left\{ g_s z_{\eta,\zeta} | \eta \in A, d(z,(\eta,\zeta)) < r, s \in [-\frac{r}{2}, \frac{r}{2}] \right\}$$

and

$$K^-(z,r,A)=\left\{g_s z_{\eta,\zeta}|\zeta\in A, d(z,(\eta,\zeta))<r, s\in[-\frac{r}{2},\frac{r}{2}]\right\}$$

Let

$$K(z,r)=K^+(z,r,PMF)=K^-(z,r,PMF)$$

Note that $\pi(K(z,r))\subseteq B(z,\frac{3r}{2})$. For $a,b\in Teich(S)$ with $d(a,b)>2r$ let

$$\Theta_r^+(a,b)=\bigcup_{w\in B(a,r)} pr_w(B(b,r))$$

and

$$\Theta_r^-(a,b)=\bigcap_{w\in B(a,r)} pr_w(B(b,r))$$

Note, as $a\to\eta\in UE$ we have $\Theta_r^{\pm}(a,b)$ converging to $\Theta_r^{\pm}(\eta,b)=\Theta_r(\eta,b)=pr_\eta(B(b,r))$.

Let $L_r(a,b)\subseteq PMF\times PMF$ denote the pairs (η,ζ) such that the geodesic defined by them passes first through $B(a,r)$ and then $B(b,r)$.

It follows immediately from the definitions that

$$L_r(a,b)\subseteq\Theta_r^+(b,a)\times\Theta_r^+(a,b).$$

Proposition 12.2. *There exists an $r_0>0$ such that for all $0<r<r_0$ and all $h>0$ the following holds. For each $\epsilon>0$ and $\epsilon'>0$ there exists an $R_0>0$ such that for every $a,b\in Teich(S)$ with the segment $[a,b]\subset Teich_\epsilon(S)$ and $d(a,b)>R_0$ such that $[a,b]$ spends at least half the time in $Q_{\epsilon'}$ we have that*

$$\Theta_r^-(b,a)\times\Theta_r^-(a,b)\subset L_{r+h}(a,b)$$

The proof of this proposition depends on the following lemmata from [5], which say that geodesic segments that spend enough time in the thick part of stratum behave like geodesics in a CAT(-1) space.

Lemma 12.3 ([5], Lemma 5.3)**.** *Suppose $K\subseteq M_g$ is compact. Given $1>\beta>0$ there exists a $\rho_0>0$ (depending only on K and β) with the following property. Given $t>0$ and $\rho>0$ there exists an $L_0=L_0(K,t,\rho,\beta)$ such that if $X,p_0\in Teich(S)$ lie above K, $d_T(p_0,p_1)<\rho_0$, $d_T(X,p_1)=L>L_0$, and*

$$|\{s\in[0,L]|l_{min}(g_s(q_{X,p_0}))\geq t\}|>\beta L$$

then

$$d_E(q,q_{X,p_0})<\rho$$

where d_E denotes the Euclidean norm and q is the unique quadratic differential in $W^{uu}(q_{X,p_0})\cap W^s(q_{X,p_1})$.

By the equivalence of the Euclidean and Teichmüller metrics over compact subsets of M_g, we also have

Lemma 12.4. *Suppose* $K \subseteq M_g$ *is compact. Given* $1 > \beta > 0$ *there exists a* $\rho_0 > 0$ *(depending only on* K *and* β*) with the following property. Given* $t > 0$ *and* $\rho > 0$ *there exists an* $L_0 = L_0(K, t, \rho, \beta)$ *such that if* $X, p_0 \in Teich(S)$ *lie above* K*,* $d_T(p_0, p_1) < \rho_0$*,* $d_T(X, p_1) = L > L_0$*, and*

$$|\{s \in [0, L] | l_{min}(g_s(q_{X,p_0})) \geq t\}| > \beta L$$

then

$$d_T(\pi(q), X) < \rho$$

where d_E *denotes the Euclidean norm and* q *is the unique quadratic differential in* $W^{uu}(q_{X,p_0}) \cap W^s(q_{X,p_1})$*.*

Lemma 12.5 ([5], Lemma 5.4)**.** *Suppose* $K \subset M_g$ *is compact. Given* $s > 0$*, there exists constants* L_0 *depending on* s *and* K*, and* c_0 *depending only on* K *with the following property. If* $\gamma : [0, L] \to Q^1(S)$ *is a geodesic segment (parametrized by arclength) with endpoints above* K *and* $L > L_0$*,* $\widehat{\gamma} : [0, L'] \to Q^1(S)$ *is the geodesic segment connecting* $p_1, p_2 \in Teich(S)$ *such that* $d_T(p_1, \pi(\gamma(0))) < c_0$*,* $d_T(p_2, \pi(\gamma(L))) < c_0$ *and*

$$|\{s \in [0, L] | l_{min}(\gamma(t)) \geq s\}| > \frac{L}{2}$$

then

$$|\{s \in [0, L'] | l_{min}(\widehat{\gamma}(t)) \geq s/4\}| > \frac{L'}{3}$$

Proof of Proposition 12.2. Let $K = Teich_\epsilon(S)/Mod(S)$. Let L_1 (depending on K and ϵ') and c_0 (depending on K) be the L_0 in Lemma 13.4 corresponding to $s = \epsilon'$ and $K = Teich_\epsilon(S)/Mod(S)$. Then, if $(x, y) \subseteq Teich_\epsilon(S)$ and $d_T(x, y) = L > L_0$ with

$$|\{s \in [0, L] | l_{min}(g_t q_{x,y}) \geq \epsilon'\}| > \frac{L}{2}$$

it follows from Lemma 12.5 that for every $p_1 \in B_{c_0}(x)$ and $p_2 \in B_{c_0}(y)$ we have that

$$|\{s \in [0, L] | l_{min}(g_t q_{x,y}) \geq \epsilon'/4\}| > \frac{L}{3}$$

Now let ρ_0 be as in Lemma 12.3, corresponding to $K = Teich_\epsilon(S)/Mod(S)$ and $\beta = 1/3$. Let $r_0 = min\{c_0, \rho_0\}$. Suppose $r < r_0$ and $h > 0$ arbitrary. Let L_2 be the L_0 in Lemma 12.4 corresponding to $K = Teich_\epsilon(S)/Mod(S)$ and $\beta = 1/3$, $t = \epsilon'/4$, and $\rho = h$. Let $R_0 > max\{L_1 + 2r_0, L_2 + 2r_0\}$. Suppose $a, b \in Teich(S)$ with the segment



Figure 1. proof of Proposition 12.2

$[a,b] \subset Teich_\epsilon(S)$ and $d(a,b) > R_0$ such that $[a,b]$ spends at least half the time outside of $Q_{\epsilon'}$ and $\eta \in \Theta_r^-(b,a)$, $\zeta \in \Theta_r^-(a,b)$. Let $b' \in B_r(b)$ be a point on the geodesic containing $[a,\zeta)$ and $a' \in B_r(a)$ a point on the geodesic containing $[b',\eta)$. We will apply Lemma 12.4 with $X = b'$, $p_0 = a'$, $p_1 = a$ to obtain that the quadratic differential $q \in W^{uu}(q_{b',\eta}) \cap W^s(q_{b',\zeta})$ satisfies $d_T(\pi(q), b') < h$, so $\pi(q) \in B_{r+h}(b)$. By definition $\pi(q) \in (\eta,\zeta)$ so (η,ζ) intersects $B_{r+h}(b)$. Similarly, (η,ζ) intersects $B_{r+h}(a)$ completing the proof. □

We now continue with the proof of Proposition 12.1. From now on, fix $r \in [0, \frac{c}{120\delta}]$ smaller than the r_0 in Proposition 12.2 such that

$$\nu_x(\partial\Theta_r(\eta_0,x)) = \nu_y(\partial\Theta_r(\zeta_0,y)) = 0$$

(this last condition only excludes countably many values of r). Since $\eta_0^* \in \Theta_r(\eta_0,x)$ and $\zeta_0^* \in \Theta_r(\zeta_0,y)$ and the conformal densities have full support on $\Lambda(G)$ we have that

$$\nu_x(\Theta_r(\eta_0,x)\nu_y(\Theta_r(\eta_0,y) > 0$$

Now, fix $h > 0$ such that

$$\nu_x(\Theta_{r-h}(\eta_0,x)) \geq e^{-c/120}\nu_x(\Theta_r(\eta_0,x))$$

and

$$\nu_y(\Theta_{r-h}(\zeta_0, y)) \geq e^{-c/120}\nu_y(\Theta_r(\zeta_0, y))$$

and also

$$\nu_x(\partial\Theta_{r-h}(\eta_0, x)) = \nu_y(\partial\Theta_{r-h}(\zeta_0, y)) = 0$$

Let $\widehat{V}$ and $\widehat{W}$ be open neighborhoods in $Teich(S) \cup PMF$ of η_0 and ζ_0 respectively such that for all $(a, b) \in (\widehat{V} \times \widehat{W})$ with $a, b \in N_D WH(G) \cup \Lambda(G)$ with $D = d(x, WH(G)) + d(y, WH(G)) + 1$ we have

$$e^{-c/120}\nu_x(\Theta_r(\eta_0, x)) \leq \nu_x(\Theta_r^{\pm}(a, x)) \leq e^{c/120}\nu_x(\Theta_r(\eta_0, x))$$

and

$$e^{-c/120}\nu_y(\Theta_r(\zeta_0, y)) \leq \nu_y(\Theta_r^{\pm}(b, y)) \leq e^{c/120}\nu_y(\Theta_r(\zeta_0, y))$$

$$e^{-c/120}\nu_x(\Theta_{r-h}(\eta_0, x)) \leq \nu_x(\Theta_{r-h}^{\pm}(a, x)) \leq e^{c/120}\nu_x(\Theta_{r-h}(\eta_0, x))$$

and

$$e^{-c/120}\nu_y(\Theta_{r-h}(\zeta_0, y)) \leq \nu_y(\Theta_{r-h}^{\pm}(b, y)) \leq e^{c/120}\nu_y(\Theta_{r-h}(\zeta_0, y))$$

It then follows that

$$e^{-c/60}\nu_x(\Theta_r(\eta_0, x)) \leq \nu_x(\Theta_{r-h}^{\pm}(a, x)) \leq \nu_x(\Theta_r^{\pm}(a, x)) \leq e^{c/60}\nu_x(\Theta_r(\eta_0, x))$$

and

$$e^{-c/60}\nu_y(\Theta_r(\zeta_0, y)) \leq \nu_y(\Theta_{r-h}^{\pm}(b, y)) \leq \nu_y(\Theta_r^{\pm}(b, y)) \leq e^{c/120}\nu_y(\Theta_r(\zeta_0, y))$$

Let V and W be open neighborhoods of η_0 and ζ_0 respectively in PMF such that $\overline{V} \subseteq \widehat{V} \cap PMF$ and $\overline{W} \subseteq \widehat{W} \cap PMF$. Consider open subsets $A \subseteq V$ and $B \subseteq W$. Let

$$K^+ = K^+(x, r, A)$$

and

$$K^+ = K^+(y, r, B)$$

We will estimate as $T \to \infty$ the quantity

$$\int_0^T e^{\delta t} \sum_{\gamma \in G} \mu(K^+ \cap g_{-t}\gamma K^-) dt$$

From the definitions, it follows that for $\gamma \in G$ and for $d(x, \gamma y) > 2r$ we have

$$\mu(K^+ \cap g_{-t}\gamma K^-) dt = \int_{L_r(x,\gamma y) \cap (\gamma B \times A)} e^{\delta \rho_x(\eta,\zeta)} d\nu_x(\eta) d\nu_y(\zeta) \int_{-r/2}^{r/2} \chi_{K(\gamma y, r)}(g_{t+s} x_{\eta,\zeta}) ds$$

We first find an upper bound for

$$\int_0^{T-3r} e^{\delta t}\sum_{\gamma\in G}\mu(K^+\cap g_{-t}\gamma K^-)dt$$

First, note that for $(\eta,\zeta)\in L_r(x,y)$ we have

$$\rho_x(\eta,\zeta)\le 2r\le c/30$$

Now, suppose $L_r(x,\gamma y)\cap(\gamma B\times A)$ is nonempty. Then from the definitions it follows that $\gamma y\in C_r^+(x,A)\subseteq C_1^+(x,A)$. Since γ is an isometry, it also holds that $L_r(\gamma^{-1}x,y)\cap(B\times\gamma^{-1}A)$ is nonempty and thus $\gamma^{-1}x\in C_1^+(y,b)$.

Note that for $(\eta,\zeta)\subseteq L_r(x,y)$, $|s|<r/2$ and $T>0$ we have

$$\int_0^{T-3r} e^{\delta t}\chi_{K(\gamma y,r)}(g_{t+s}x_{\eta,\zeta})dt\le e^{\delta(3r)}re^{\delta d(x,\gamma y)}\le e^{c/20}re^{\delta d(x,\gamma y)}$$

and moreover is zero whenever $d(x,\gamma y)>T$. Using the fact that $L_r(a,b)\subseteq\Theta_r^+(b,a)\times\Theta_r^+(a,b)$ it follows that

$$\int_0^{T-3r} e^{\delta t}\sum_{\gamma\in G}\mu(K^+\cap g_{-t}\gamma K^-)dt\le e^{c/12}r^2\sum\nu_x(\Theta_r^+(\gamma y,x))\nu_y(\Theta_r^+(x,\gamma y))$$

where the sum is taken over all $\gamma\in G$ such that $(x,\gamma y)\le T$ and

$$(\gamma y,\gamma^{-1}x)\in[C_1^+(x,A)\times C_1^+(y,B)]$$

By Corollary 7.4 we see that he set

$$[C_1^+(x,A)\cap N_DWH(G)\times C_1^+(y,B)\cap N_DWH(G)]\setminus[\widehat{V}\times\widehat{W}]$$

has compact closure in $Teich(S)$ for any $D>0$. If $x,y\in N_DWH(G)$ their G orbits are also contained in $N_DWH(G)$. Thus, by the discreteness of the G action on $Teich(S)$, for some constant c_1 that does not depend on T

$$\int_0^{T-3r} e^{\delta t}\sum_{\gamma\in G}\mu(K^+\cap g_{-t}\gamma K^-)dt\le e^{c/12}r^2\sum\nu_x(\Theta_r^+(\gamma y,x))\nu_y(\Theta_r^+(x,\gamma y))-c_1$$

for all $T>0$ where the sum is taken over all $\gamma\in G$ with $(x,\gamma y)\le T$ and $(\gamma y,\gamma^{-1}x)\in[C_1^+(x,A)\times C_1^+(y,B)]\cap[\widehat{V}\times\widehat{W}]$.

Note, by the triangle inequality we have for $\eta\in\Theta_r^+(x,\gamma y))$

$$d(x,\gamma y)-4r\le\beta_\eta(x,\gamma y)\le d(x,\gamma y)$$

and by the conformality of ν it follows that

$$\nu_y(\Theta_r^+(\gamma^{-1}x,y))=\nu_{\gamma y}(\Theta_r^+(x,\gamma y))\le\nu_x(\Theta_r^+(x,\gamma y))e^{\delta d(x,\gamma y)}\le e^{4r}\nu_{\gamma y}(\Theta_r^+(x,\gamma y))$$

Since $4\delta r < c/15$ it follows that when

$$(\gamma y, \gamma^{-1}x) \subseteq \widehat{V} \times \widehat{W}$$

we have that

$$\begin{aligned}\nu_x(\Theta_r^+(\gamma y, x))\nu_x(\Theta_r^+(x, \gamma y)) &\leq e^{c/15}\nu_x(\Theta_r^+(\gamma y, x))\nu_y(\Theta_r^+(\gamma^{-1}x, y)) \\ &\leq e^{c/10}\nu_x(\Theta_r(\eta_0, x))\nu_y(\Theta_r(\zeta_0, y))\end{aligned}$$

Thus we obtain

$$\int_0^{T-3r} e^{\delta t}\sum_{\gamma\in G}\mu(K^+\cap g_{-t}\gamma K^-)dt \leq e^{c/6}r^2|G^+(T,A,B)|\nu_x(\Theta_r(\eta_0, x))\nu_y(\Theta_r(\zeta_0, y))+c_1$$

where c_1 is independent of T and $G^+(T, A, B)$ is the set of all $\gamma \in G$ such that

$$(x, \gamma y) \leq T$$

and

$$(\gamma y, \gamma^{-1}x) \in [C_1^+(x, A) \times C_1^+(y, B)] \cap [\widehat{V} \times \widehat{W}]$$

In a similar but more annoying manner, we will obtain a lower bound for

$$\int_0^{T+3r} e^{\delta t}\sum_{\gamma\in G}\mu(K^+ \cap g_{-t}\gamma K^-)dt$$

First, note that for $\eta \in \Theta^-_{r-h}(a, b)$ we have

$$d(a, b) - 2r \leq \beta_\eta(a, b) \leq d(a, b)$$

and thus similarly to above we have

$$\nu_y(\Theta^-_{r-h}(\gamma^{-1}x, y)) = \nu_{\gamma y}(\Theta^-_{r-h}(x, \gamma y)) \leq \nu_x(\Theta^-_{r-h}(x, \gamma y))e^{\delta d(x,\gamma y)} \leq e^{2r}\nu_{\gamma y}(\Theta^-_{r-h}(x, \gamma y))$$

Since $2\delta r < c/30$ it follows that when

$$(\gamma y, \gamma^{-1}x) \subseteq \widehat{V} \times \widehat{W}$$

we have that

$$\begin{aligned}\nu_x(\Theta^-_{r-h}(x, \gamma y))\nu_x(\Theta^-_{r-h}(\gamma y, x)) &\geq e^{-\delta d(x,\gamma y)}\nu_x(\Theta^-_{r-h}(x, \gamma y))\nu_y(\Theta^-_{r-h}(x, \gamma y)) \\ &\geq e^{-c/30}\nu_x(\Theta_r(\eta_0, x))\nu_y(\Theta_r(\zeta_0, y))\end{aligned}$$

Now note, if $(\gamma y, \gamma^{-1}x) \in C_1^-(x, A) \times C_1^-(y, B)$ then by definition

$A \supset \Theta_r^-(x, \gamma y)$ and $B \supset \Theta_r^-(y, \gamma^{-1}x)$, whence $\gamma B \supset \Theta_r^-(\gamma y, x)$. Note for $(\eta, \zeta) \in L_r(x, y)$, $|s| < r/2$, $T > 0$ and $3r \leq d(x, \gamma y) \leq T$ we have

$$\int_0^{T+3r} e^{\delta t}\chi_{K(\gamma y,r)}(g_{t+s}x_{\eta,\zeta})dt \geq e^{-3\delta r}re^{\delta d(x,\gamma y)} \geq e^{-c/20}re^{\delta d(x,\gamma y)}$$

Now fix an $\epsilon' > 0$ with $\mu^{BMS}(Q_{\epsilon'}(S)/G) < 1/3$ and consider $\gamma \in G$ such that $[x, \gamma y]$ and $[\gamma y, x]$ both spend less than half time in $Q_{\epsilon'}$. By Proposition 12.6 and the discreteness of the action of G, for all but finitely many such γ we have that

$$\Theta^-_{r-h}(\gamma y, x) \times \Theta^-_{r-h}(x, \gamma y) \subset L_r(x, \gamma y)$$

Note for $(\eta, \zeta) \in L_r(x, y), |s| < r/2$, $T > 0$ and $3r \leq d(x, \gamma y) \leq T$ we have

$$\int_0^{T+3r} e^{\delta t} \chi_{K(\gamma y, r)}(g_{t+s} x_{\eta,\zeta}) dt \geq e^{-c/20} r e^{\delta d(x,\gamma y)}$$

Thus we have that

$$\int_0^{T+3r} e^{\delta t} \sum_{\gamma \in G} \mu(K^+ \cap g_{-t}\gamma K^-) dt \geq e^{-c/20} r^2 \sum_{\gamma \in G(T,A,B)} \nu_x(\Theta^-_{r-h}(\gamma y, x)) \nu_x(\Theta^-_{r-h}(x, \gamma y)) e^{\delta d(x,\gamma y)} \geq$$

$$e^{-c/12} r^2 |G^-(T, A, B) \setminus G^-(T, \epsilon', A, B)| \nu_x(\Theta_r(\eta_0, x)) \nu_y(\Theta_r(\zeta_0, y)) - c_2$$

where $G^-(T, A, B)$ is the set of all $\gamma \in G$ such that $(x, \gamma y) \leq T$ and

$$(\gamma y, \gamma^{-1} x) \in [C_1^+(x, A) \times C_1^+(y, B)] \cap [\widehat{V} \times \widehat{W}]$$

and $G^-(T, \epsilon', A, B)$ is the set of all $\gamma \in G(T, A, B)$ such that the segment $(x, \gamma y)$ spends at least half the time in the ϵ' thin part of the principal stratum, and c_2 does not depend on T.

By mixing it follows that for all t large enough we have

$$e^{-c/60} \mu(K^+) \mu(K^-) \leq ||\mu|| \sum_{\gamma \in G} \mu(K^+ \cap g^{-t} \gamma K^-) \leq e^{c/60} \mu(K^+) \mu(K^-)$$

Note, that by definition

$$\mu(K^+) = r \int_{\eta \in A} \int_{\zeta \in \Theta_r(\eta, x)} e^{\delta \rho_x(\eta,\zeta)} d\nu_x(\zeta) d\nu_x(\eta)$$

Since

$$0 \leq \rho_x(\eta, \zeta) \leq 2r$$

for $\zeta \in \Theta_r(\eta, x)$ and since $A \subseteq V$ we obtain that

$$e^{-c/60} r \nu_x(A) \nu_x(\Theta_r(\eta_0, x)) \leq \mu(K^+) \leq e^{c/20} r \nu_x(A) \nu_x(\Theta_r(\eta_0, x))$$

and similarly

$$e^{-c/60} r \nu_y(A) \nu_y(\Theta_r(\zeta_0, y)) \leq \mu(K^-) \leq e^{c/20} r \nu_y(A) \nu_y(\Theta_r(\zeta_0, y))$$

It follows that there exists a constant c_2 independent of T such that

$$\delta ||\mu|| \int_0^{T-3r} e^{\delta t} \sum_{\gamma \in G} \mu(K^+ \cap g_{-t} \gamma K^-) dt \geq e^{-c/2} e^{\delta T} M \nu_x(A) \nu_y(B) - c_2$$

and

$$\delta||\mu||\int_0^{T+3r} e^{\delta t}\sum_{\gamma\in G}\mu(K^+\cap g_{-t}\gamma K^-)dt \leq e^{c/2}e^{\delta T}M\nu_x(A)\nu_y(B)+c_2$$

where $M = r^2\nu_x(\Theta_r(\eta_0,x))\nu_y(\Theta_r(\zeta_0,y))$.

Thus, it follows that

$$e^{-c/2}\nu_x(A)\nu_x(B) \leq e^{c/3}e^{-\delta T}|G^+(T,A,B)|$$

and

$$e^{c/2}\nu_x(A)\nu_x(B) \geq e^{-c/6}e^{-\delta T}|G^-(T,A,B)\setminus G^-(T,\epsilon' A,B)|$$

for large enough T. Furthermore, by Theorem 11.4, if ϵ' is chosen small enough,

$$\limsup_{T\to\infty} |G^-(T,\epsilon' A,B)|/e^{\delta t} < e^{c/12}$$

so that

$$e^c\nu_x(A)\nu_x(B) \geq e^{-\delta T}|G^-(T,A,B)|$$

for all large enough T.

This completes the proof of Proposition 12.1. □

Lemma 12.6. *Let $x, y \in Teich(S)$ and $c > 0$. For each*

$$(\eta_0,\zeta_0)\in PMF\times PMF$$

there exists an $r > 0$ and neighborhoods V and W of η_0 and ζ_0 in PMF respectively such that for all borel $A \subseteq V$ and $B \subseteq W$, with nonempty interior:

$$\limsup_{t\to\infty} \nu^t_{x,y}(C^-_r(x,A)\times C^-_r(y,B)) \leq e^c\nu_x(A)\nu_y(B)$$

and

$$\liminf_{t\to\infty} \nu^t_{x,y}(C^+_r(x,A)\times C^+_r(y,B)) \geq e^{-c}\nu_x(A)\nu_y(B)$$

Proof. If η_0 is not in $\Lambda(G)$ then we can choose a neighborhood U of η_0 in PMF with $\nu_x(U) = 0$ and $W = PMF$ so that both sides of the desired equation are 0 by Corollary 9.7. Similarly if ζ_0 is not in $\Lambda(G)$. Assume therefore that $\eta_0,\zeta_0 \in \Lambda(G)$. Let $\lambda_0 \in \Lambda(G)$ and $x_0 \in (\eta_0,\lambda_0)$, $y_0 \in (\zeta_0,\lambda_0)$. Let V_0, W_0 be open neighborhoods of η_0 and ζ_0 in PMF respectively such that for all open $A \subseteq V_0$ and $B \subseteq W_0$, we have as $T\to\infty$ that

$$\limsup \nu^T_{x_0,y_0}(C^-_1(x_0,A)\times C^-_r(y_0,B)) \leq e^{c/3}\nu_{x_0}(A)\nu_{y_0}(B)$$

and

$$\liminf \nu^T_{x_0,y_0}(C^+_r(x_0,A)\times C^+_r(y_0,B)) \geq e^{-c/3}\nu_{x_0}(A)\nu_{y_0}(B)$$

Let $\widehat{V_0}$ and $\widehat{W_0}$ be neighborhoods in $Teich(S) \cup PMF$ of η_0 and ζ_0 respectively, whose intersection with PMF are respectively contained in V_0 and W_0 and such that for all

$$a \in \widehat{V_0} \cap N_D WH(G), b \in \widehat{W_0} \cap N_D WH(G)$$

we have

$$|d(x_0, a) - d(x, a) - \beta_{\eta_0}(x_0, x)| \leq \frac{c}{6\delta}$$

$$|d(y_0, b) - d(y, b) - \beta_{\eta_0}(y_0, y)| \leq \frac{c}{6\delta}$$

and for all

$$\eta \in \widehat{V_0} \cap \Lambda(G)$$

$$\zeta \in \widehat{W_0} \cap \Lambda(G)$$

we have

$$|\beta_{\eta}(x_0, x) - \beta_{\eta_0}(x_0, x)| \leq \frac{c}{6\delta}$$

and

$$|\beta_{\eta}(x_0, x) - \beta_{\eta_0}(x_0, x)| \leq \frac{c}{6\delta}.$$

Let V and W be open neighborhoods in PMF of η_0 and ζ_0 respectively, with $\overline{V} \subseteq \widehat{V_0} \cap PMF$ and $\overline{W} \subseteq \widehat{W_0} \cap PMF$ and let

$$r = 1 + d(x, x_0) + d(y, y_0).$$

Consider $A \subseteq V$ and $B \subseteq W$.

Note, if $(\gamma y, \gamma^{-1} x) \in C_r^-(x, A) \times C_r^-(y, B)$ one easily checks that $(\gamma y_0, \gamma^{-1} x_0) \in C_1^-(x_0, A) \times C_1^-(y_0, B)$ by the choice of r. Next, note that if $d(x, \gamma y) \leq t$ and $(\gamma y, \gamma^{-1} x) \in \widehat{V_{-r}} \times \widehat{W_0}$ where $\widehat{V_{-r}}$ denotes the set of points whose r neighborhood is contained in $\widehat{V_0}$, then $\gamma y_0 \in \widehat{V_0}$ and $\gamma^{-1} x \in \widehat{W_0}$ which implies that

$$d(x_0, \gamma y_0) \leq d(x, \gamma y_0) + \beta_{\eta_0}(x_0, x) + \frac{c}{6\delta} = d(y_0, \gamma^{-1} x) + \beta_{\eta_0}(x_0, x) + \frac{c}{6\delta}$$

$$\leq d(y, \gamma^{-1} x) + \beta_{\zeta_0}(y_0, y) + \beta_{\eta_0}(x_0, x) + \frac{c}{3\delta} \leq t + \beta_{\eta_0}(x_0, x) + \beta_{\zeta_0}(y_0, y) + \frac{c}{3\delta}$$

From Corollary 7.4 we obtain

Lemma 12.7.

$$[C_r^-(x, A) \cap N_D WH(G) \times C_r^-(y, B) \cap N_D WH(G)] \setminus [\widehat{V_{-r}} \times \widehat{W_0}]$$

is relatively compact in $Teich(S)$.

Thus $[C_r^-(x,A)\times C_r^-(y,B)]\setminus[\widehat{V_{-r}}\times\widehat{W_0}]$ contains only finitely many points $(\gamma x,\gamma^{-1}y)$, $\gamma\in G$

From this, we deduce that

$$\limsup_{t\to\infty}\nu_{x,y}^t(C_r^-(x,A)\times C_r^-(y,B))\leq$$

$$e^{c/3}e^{\delta\beta_{\eta_0}(x_0,x)+\delta\beta_{\zeta_0}(y_0,y)}\limsup\nu_{x_0,y_0}^{t+\beta_{\eta_0}(x_0,x)+\beta_{\zeta_0}(y_0,y)+\frac{c}{3\delta}}(C_1^-(x_0,A)\times C_1^-(y_0,B))$$

and thus by Prop 12.1,

$$\limsup\nu_{x,y}^t(C_r^-(x,A)\times C_r^-(y,B))\leq e^{2c/3}e^{\delta\beta_{\eta_0}(x_0,x)+\delta\beta_{\zeta_0}(y_0,y)}\nu_{x_0}(A)\nu_{y_0}(B).$$

Since $e^{\delta\beta_{\eta_0}(x_0,x)}\nu_{x_0}\leq e^{c/6}\nu_x$ when restricted to V and $e^{\delta\beta_{\eta_0}(y_0,y)}\nu_{y_0}\leq e^{c/6}\nu_y$ when restricted to W, we obtain that

$$\limsup\nu_{x,y}^t(C_r^-(x.A)\times C_r^-(y.B))\leq e^c\nu_x(A)\nu_y(B)$$

The reverse estimate is proved similarly. □

Theorem 12.8. *For $x,y\in Teich(S)$ $\nu_{x,y}^t$ converges weakly to $\nu_x\times\nu_y$ as $t\to\infty$*

Proof. Let $c>0$. For each $(\eta_0,\zeta_0)\in\Lambda(G)\times\Lambda(G)$ take neighborhoods $V_{(\eta_0,\zeta_0)}$ and $W_{(\eta_0,\zeta_0)}$ of η_0 and ζ_0 respectively such that the conclusion of Lemma 12.6 holds for c. By compactness finitely many of the $V\times W$ cover $PMF\times PMF$, say $V_i\times W_i$, $i=1,...,n$. Let $\widehat{V_i}$ and $\widehat{W_i}$ be open subsets of $Teich(S)\cup PMF$ such that $V_i=\widehat{V_i}\cap PMF$ and $W_i=\widehat{W_i}\cap PMF$. Let $\widehat{A}$ and $\widehat{B}$ be borel subsets of $Teich(S)\cup PMF$ with

$$\overline{\widehat{A}}\subset\widehat{V_i}$$

$$\overline{\widehat{B}}\subset\widehat{W_i}$$

and

$$(\nu_x\otimes\nu_y)(\partial(\widehat{A}\times\widehat{B}))=0$$

ie

$$\nu_x(\overline{\widehat{A}}))\nu_y(\partial\widehat{B})=\nu_x(\overline{\widehat{B}}))\times\nu_y(\partial\widehat{A})=0$$

Let $\alpha>0$. Let $A^+,B^+\subset PMF$ be open and $A^-,B^-\subset PMF$ compact with A^-,B^- either being empty or having nonempty interior such that

$$A^-\subset\widehat{A}^o\cap PMF\subset\overline{\widehat{A}}\cap PMF\subset A^+\subset V_i$$

$$B^-\subset\widehat{B}^o\cap PMF\subset\overline{\widehat{B}}\cap PMF\subset B^+\subset W_i$$

$$\nu_x(\widehat{A}^o\setminus A^-)<\alpha,\nu_x(A^+\setminus\overline{\widehat{A}})<\alpha,\nu_x(\widehat{B}^o\setminus B^-)<\alpha,\nu_x(A^+\setminus\overline{\widehat{A}})<\alpha$$

Let $D > d(x, WH(G)) + d(y, WH(G))$ so that the G orbits of x, y are contained in $N_D WH(G)$. By Corollary 7.4 the sets

$$[\overline{\widehat{A}} \cap N_D WH(G) \times \overline{\widehat{B}} \cap N_D WH(G)] \setminus [C_r^-(x, A^+) \times C_r^-(y, B^+)]$$

and

$$[C_r^+(x, A^-) \cap N_D WH(G) \times C_r^+(y, B^-) \cap N_D WH(G)] \setminus [\widehat{A}^o \times \widehat{B}^o]$$

are relatively compact in $Teich(S) \times Teich(S)$. Thus, by Lemma 12.6 we have

$$\limsup \nu^t_{x,y}(\widehat{A}\times\widehat{B}) \leq \limsup \nu^t_{x.y}(C_r^-(x, A^+)\times C_r^-(y, B^+)) \leq e^c \nu_x(A^+)\nu_y(B^+) \leq$$

$$e^c \nu_x(\overline{\widehat{A}})\nu_y(\overline{\widehat{B}}) + \alpha e^c(||\nu_x|| + ||\nu_y||) = e^c \nu_x(\widehat{A})\nu_y(\widehat{B}) + \alpha e^c(||\nu_x|| + ||\nu_y||)$$

Since $\alpha > 0$ can be chosen arbitrarily small we obtain

$$\limsup \nu^t_{x,y}(\widehat{A} \times \widehat{B}) \leq e^c \nu_x(\overline{A})\nu_y(\overline{B})$$

Similarly we obtain the reverse estimate

$$\liminf \nu^t_{x,y}(\widehat{A} \times \widehat{B}) \geq e^{-c} \nu_x(\overline{A})\nu_y(\overline{B})$$

Indeed,

$$\liminf \nu^t_{x,y}(\widehat{A} \times \widehat{B}) \geq \liminf \nu^t_{x.y}(C_r^+(x, A^-) \times C_r^+(y, B^-)) \geq$$

$$e^{-c}\nu_x(\widehat{A}^o)\nu_y(\widehat{B}^o) - \alpha e^c(||\nu_x|| + ||\nu_y||) = e^{-c}\nu_x(\widehat{A})\nu_y(\widehat{B}) - \alpha e^c(||\nu_x|| + ||\nu_y||)$$

for any $\alpha > 0$. Thus, for ϕ a continuous function supported on $\widehat{V} \times \widehat{W}$ we have

$$e^{-c}\int \phi d\nu_x \otimes d\nu_y \leq \liminf \int \phi d\nu^t_{x,y} \leq \liminf \int \phi d\nu^t_{x,y} \leq e^{-c}\int \phi d\nu_x \otimes d\nu_y$$

Furthermore the complement O of $\bigcup_{i=1}^n V_i \times W_i$ in $Teich(S) \cup PMF$ is a compact subset of $Teich(S)$, so for any function supported on O we have

$$\int \phi d\nu_x \otimes d\nu_y = 0$$

and

$$\lim_{t\to\infty} \int \phi d\nu^t_{x,y} = 0$$

By choosing a partition of unity subordinate to the cover $O, \widehat{V_i} \times \widehat{W_i}$ we obtain that

$$e^{-c}\int \phi d\nu_x \otimes d\nu_y \leq \liminf \int \phi d\nu^t_{x,y} \leq \liminf \int \phi d\nu^t_{x,y} \leq e^{-c}\int \phi d\nu_x \otimes d\nu_y$$

for any continuous ϕ on $(Teich(S) \cup PMF) \times (Teich(S) \cup PMF)$ and letting $c \to 0$ yields the desired result.

□

Theorem 1.1 follows.

Theorem 12.9. *For all $x, y \in Teich(S)$ we have*

$$\lim_{R\to\infty} |B_R(x) \cap Gy|e^{-\delta R} = \frac{||\nu_x||||\nu_y||}{\delta||\mu^{BMS}||}$$

13. Counting Closed Geodesics

In this section we prove Theorem 1.2. Denote by G_h the set of pseudo-Anosov elements of G. Denote by G_{hp} the set of primitive pseudo-Anosov elements of G. Let $\Omega(l)$ be the set of closed primitive geodesics on $Teich(S)/G$ of length at most R. For $g \in \Omega(l)$ let D_g be the Lebesgue measure on g normalized to unit mass.

We will prove:

Theorem 13.1.

$$\lim_{t\to\infty} \delta t e^{-\delta t} \sum_{g\in\Omega(t)} D_g = ||\mu||^{-1}\mu$$

Theorem 1.1 then follows.

Proof of Theorem 13.1. Let $x \in Teich(S)$. Denote by $V(x,r) \subset Teich(S)^2 \cup PMF^2$ the set of pairs (a,b) such that $[a,b] \cap B(x,r) \neq \emptyset$. Recall the measure $\widetilde{\mu}$ on $\Lambda(G) \times \Lambda(G)$ by

$$d\tilde{\mu}(\eta,\zeta) = \exp(\delta(G)\rho_x(\eta,\zeta))d\nu_x(\eta)d\nu_x(\zeta)$$

and let

$$d\nu^t_{x,1} = d\nu^t_{x,x} = \delta||\mu||e^{-\delta t} \sum_{\gamma\in G, d(x,\gamma y)\le t} D_{\gamma x} \otimes D_{\gamma^{-1}x}$$

$$d\nu^t_{x,2} = \delta||\mu||e^{-\delta t} \sum_{\gamma\in G_h, d(x,\gamma y)\le t} D_{\gamma x} \otimes D_{\gamma^{-1}x}$$

$$d\nu^t_{x,3} = \delta||\mu||e^{-\delta t} \sum_{\gamma\in G_h, d(x,\gamma y)\le t} D_{\gamma^+} \otimes D_{\gamma^-}$$

where $\gamma^\pm \in PMF$ denote the stable and unstable laminations of γ. Note that for $D = d(x, WH(G))$ we have $\nu^t_{x,i}$ and $\nu_x \times \nu_x$ all supported on $(N_D WH(G) \cup \Lambda(G))^2$ and $V(x,r) \cap (N_D WH(G) \cup \Lambda(G))^2$ is closed in $Teich(S) \cup PMF$.

Lemma 13.2. *For every $c > 0$ there exists a $t_0 = t_0(x,r,c) > 0$ such that if $\gamma \in G$ with $d(x,\gamma x) > t_0$ and $(\gamma x, \gamma^{-1}x) \in V(x,r)$ we have that γ is pseudo-Anosov and $\rho_x(\gamma^{\pm 1}x, \gamma^\pm) > c$.*

Proof. Suppose $\gamma \in G$ is a pseudo-Anosov, p a point on the axis of γ and l the unit speed parametrization of the axis starting at p in direction γ^+. Note,

$$\rho_x(\gamma x, \gamma^+) = \lim_{t\to\infty} d(x, \gamma x) + d(x, l(t)) - d(\gamma x, l(t)) =$$

$$d(x, \gamma x) + d(x, l(t)) - d(x, l(t - l(\gamma))) \geq d(x, \gamma x)$$

and similarly

$$\rho_x(\gamma^{-1}x, \gamma^-) \geq d(x, \gamma x)$$

Note, as G is convex cocompact it contains no parabolic elements. Since G is a hyperbolic group, it contains only finitely many conjugacy classes of finite order elements. Therefore, the fixed points of finite order elements of G are contained in finitely many G orbits of $Teich(S)$. Let D be the maximum distance of these orbits from $WH(G)$. Suppose now that $\gamma_n \in G$ is a sequence of finite order elements with $d(x, \gamma_n x) \to \infty$ and $(\gamma_n x, \gamma_n^{-1}x) \in V(x, r)$. Let p_n be the fixed point of γ_n. Then $x, \gamma_n x, \gamma_n^{-1}x$ all lie on the same circle C_n of radius $r_n \to \infty$ centered at p_n.

Taking a subsequence, we can assume

$$p_n \to \eta \in PMF$$

and

$$y_n = \gamma_n x \to \zeta \in PMF$$

and

$$z_n = \gamma_n^{-1} x \to \theta \in PMF$$

Then clearly $\zeta, \theta \in \Lambda(G)$.

Also, the p_n are all contained in $N_D WH(G)$ and thus $\eta \in \Lambda(G) \subset UE$. We claim $\zeta = \eta = \theta$, which would imply that for large enough n we have $(\gamma_n x, \gamma_n^{-1}x) \notin V(x, r)$, contradicting our assumption. Indeed,

$$\rho_x(\eta, \zeta) = \lim_{n\to\infty} d(x, p_n) + d(x, y_n) - d(y_n, p_n) = \lim_{n\to\infty} r_n \to \infty$$

which is impossibe if $\eta \neq \zeta$ by continuity of ρ. Similarly, $\eta = \theta$. □

For the remainder of the argument, the proof of Roblin's Theorem 5.1.1 carries through with essentially no modification. From Lemma 13.2 we obtain

Corollary 13.3. *When restricted to* $V(x, r)$ *we have* $\nu^t_{x,i} - \nu^t_{x,j} \to 0$ *for* $i, j = 1, 2, 3$.

Note, for $\eta, \zeta \in V(x,r)$ we have $0 < \rho_x(\eta,\zeta) \leq 2r$ and thus by Theorem 12.8 and Lemma 13.2 for any positive continuous ψ compactly supported on $V(x,r)$ we have

$$e^{-2\delta r}\int \psi d\widetilde{\mu} \leq \liminf \int \psi d\nu_{x,3}^t \leq \limsup \int \psi d\nu_{x,3}^t \leq \int \psi d\widetilde{\mu}$$

as $t \to \infty$. For $\gamma \in G_h$ let g_γ be the axis of γ, and denote by L_γ the arclength measure on γ. Let $l(\gamma)$ denote the translation length of γ in the Teichmüller metric. Let

$$M_x^t = \delta e^{-\delta t} \sum_{\gamma \in G_h, l(\gamma) \leq t} L_\gamma$$

and

$$M_{x,3}^t = \delta e^{-\delta t} \sum_{\gamma \in G_h, d(x,\gamma x) \leq t} L_\gamma$$

Note that

$$l(\gamma) \leq d(x,\gamma x) \leq l(\gamma) + 2d(x, g_\gamma)$$

and thus when restricted to $V(x,r)$ we have

$$M_{x,3}^t \leq M_x^t \leq e^{2\delta r} M_{x,3}^{t+2r}$$

Let $\widehat{V}(x,r) \subset Q^1(S)$ denote all quadratic differentials on geodesic segments defined by elements of $V(x,r)$. Note,

$$M_{x,3}^t = ||\mu^{BMS}||^{-1} \nu_{x,3}^t \otimes ds$$

and thus for any $\phi \in C_c^+(\widehat{V}(x,r))$ we have

$$e^{-2\delta r}||\mu||^{-1}\int \phi d\mu \leq \liminf \int \phi dM_{x,3}^t \leq \limsup \int \phi dM_{x,3}^t \leq ||\mu||^{-1}\int \phi d\mu$$

Denote by $G_{hp} \subset G$ the set of primitive hyperbolic isometries so that

$$M_x^t = \delta e^{-\delta t} \sum_{\gamma \in G_{hp}, l(\gamma) \leq t} \left\lceil \frac{t}{l(\gamma)} \right\rceil L_\gamma$$

Clearly

$$M_x^t \leq E^t := \delta t e^{-\delta t} \sum_{\gamma \in G_{hp}, l(\gamma) \leq t} \frac{1}{l(\gamma)} L_\gamma$$

Moreover, note $\left\lceil \frac{t}{l(\gamma)} \right\rceil \geq \frac{1}{l(\gamma)}$ whenever $1 \leq l(\gamma) \leq t$ and $t \geq 2$, so for any $\phi \in C_c^+(\widehat{V(x,r)})$ we have

$$\sum_{\gamma \in G_{hp}, l(\gamma) \leq t} \frac{1}{l(\gamma)} \int \phi dL_\gamma = O(e^{\delta t})$$

since $\int \phi dM_x^t$ is bounded as $t \to \infty$.

Now, if $e^{-r}t < l(\gamma) \leq t$ then

$$\left\lceil \frac{t}{l(\gamma)} \right\rceil \geq 1 \geq \frac{e^{-r}t}{l(\gamma)}$$

and so

$$\int \phi dM_x^t \geq e^{-r}\delta t e^{-\delta t} \sum_{\gamma \in G_{hp}, e^{-r}t < l(\gamma) \leq t} \frac{1}{l(\gamma)} \int \phi dL_\gamma =$$

$$e^{-r} \int \phi dE^t - e^{-r}\delta t e^{-\delta t} \sum_{\gamma \in G_{hp}, l(\gamma) \leq e^{-r}t} \frac{1}{l(\gamma)} \int \phi dL_\gamma$$

By above remarks, the second term in the above difference is bounded above by a constant multiple of

$$t e^{\delta(e^{-r}-1)t} = o(1)$$

Thus we get

$$\limsup \int \phi dM_x^t \geq e^{-r} \limsup \int \phi dE^t$$

Putting everything together we obtain as $t \to \infty$

$$e^{-2\delta r}||\mu||^{-1} \int \phi d\widetilde{\mu} \leq \limsup \int \phi dE^t \leq \limsup \int \phi dE^t \leq e^{(2\delta+1)r}||\mu||^{-1} \int \phi d\widetilde{\mu}$$

Choosing a partition of unity subordinate to the locally finite cover of $Q^1(S)$ by the $\widehat{V(x,r)}$ (where $r > 0$ is fixed) we obtain the above relation for each $\phi \in C_c^+(Q^1(S))$. Letting $r \to 0$ completes the proof. □

References

[1] Jayadev Athreya, Alexander Bufetov, Alex Eskin and Maryam Mirzakhani. Lattice Point Asymptotics and Volume Growth on Teichmller space. Preprint, arXiv:math/0610715v3, 2010, 139.

[2] Benson Farb and Lee Mosher. Convex cocompact subgroups of mapping class groups. Geom. Topol., 6:91152 (electronic), 2002.

[3] Giovanni Forni. Deviation of ergodic averages for area-preserving ows on surfaces of higher genus. Ann. of Math. (2), 155 (2002), no. 1, 1103.

[4] Moon Duchin, Thin triangles and a multiplicative ergodic theorem for Teichmller geometry. Preprint, arXiv:math.GT/0508046.

[5] Alex Eskin, Maryam Mirzakhani. Counting closed geodesics in Moduli space. Preprint, arXiv:0811.236

[6] Alex Eskin, Maryam Mirzakhani, and Kasra Rafi. Counting Closed Geodesics in Strata. Preprint,
http://www.math.ou.edu/∼rafi/research/Loop.pdf

[7] A. Fathi, F. Laudenbach, and V. Poenaru. Travaux de Thurston sur les surfaces. Societ e Mathematique de France, Paris, 1991. Seminaire Orsay, Reprint of Travaux de Thurston sur les surfaces, Soc. Math. France, Paris, 1979 Asterisque No. 66-67 (1991).

[8] Roland Gunesch. Counting closed geodesics on rank one manifolds. Preprint, http://arxiv.org/pdf/0706.2845v1.pdf

[9] Roland Gunesch. Precise volume estimates in nonpositive curvature. Preprint, www.math.uni-hamburg.de/home/gunesch/papers/volume.ps

[10] Ursula Hamenstaedt. Bowen's construction for the Teichmüller Flow. Preprint, http://arxiv.org/abs/1007.2289

[11] John Hubbard and Howard Masur, Quadratic Differentials and Foliations. Acta Math., 142 (3-4):221-274,1979.

[12] Gregory Margulis. On some aspects of the theory of Anosov systems. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2004.

[13] E. Klarreich, the boundary at Infinity of the Curve Complex and the Relative Teichmüller Space Preprint, http://www.ericaklarreich.com/curvecomplex.pdf

[14] R. P. Kent, and C. J. Leininger, Shadows of mapping class groups: capturing convex cocompactness. Geom. Funct. Anal. 18 (2008), 12701325

[15] R. P. Kent, and C. J. Leininger, Subgroups of mapping class groups from the geometrical viewpoint. In the tradition of Ahlfors-Bers. IV, 119141, Contemp. Math., 432, Amer. Math. Soc., Providence, RI, 2007

[16] Howard Masur Uniquely ergodic quadratic dierentials, Comment. Math. Helvetici Vol. 55 (1980) 255-266

[17] Howard Masur Interval exchange transformations and measured foliations, Annals of Math. 115 169-200 (1982)

[18] John McCarthy and Athanase Papadopoulos. Dynamics on Thurston's sphere of projective measured foliations. Comment. Math. Helv., 64(1):133166, 1989.

[19] Howard Masur, and Yair Minsky, Geometry of the complex of curves. I. Hyperbolicity. Inventiones Mathematicae, vol. 138 (1999), no. 1, pp. 103149.

[20] Hideki Miyachi. Teichmüller rays and the Gardiner-Masur boundary of Teichmüller space. Geom. Dedicata 137 (2008), 113141.

[21] Kasra Rafi. Hyperbolicity in Teichmüller Space. Preprint, http://www.math.ou.edu/∼rafi/research/Fellow.pdf

[22] Thomas Roblin. Ergodicity and uniform distributions in negative curvature. Mem. Soc. Math. Fr., No. 95, 2003.

[23] Yves Guivarc'h, Roman Urban, Semigroup actions on tori and stationary measures on projective spaces, Studia Math. 171 (1) (2005) 33 66.

[24] Martine Babillot. On the mixing property for hyperbolic systems. Israel J. Math. 129 (2002), 6176

[25] J.F. Quint. An Overview of Patterson Sullivan Theory. www.math.univ-paris13.fr/ quint/publications/courszurich.pdf

[26] D. Sullivan, The density at infinity of a discrete group of hyperbolic motions, Publ. Math. I.H.E.S. 50 (1979), 171-202. MR 81b:58031

[27] Chengbo Yue. The ergodic theory of discrete isometry groups on manifolds of variable negative curvature, Trans. Amer. Math. Soc. 348 (1996), no. 12, 49655005.

[28] Michel Coornaert. Sur les groups proprement discontinus d'isometries des espaces hyper- boliques au sens de Gromov, These (1990).

[29] Gerhard Knieper. On the asymptotic geometry of nonpositively curved manifolds, Geom. Funct. Anal. 7 (1997), 755782.

[30] Gabriele Link. Asymptotic Geometry and Growth of Conjugacy Classes of Nonpositively Curved Manifolds Annals of Global Analysis and Geometry. Volume 31, Number 1, 37-57.